




Figure 1. Top: Topography of Euro-Asian continent. The Caspian Sea, currently  $\sim 28\text{m}$  below sea level, shown in blue, bottom: CZCS average pigment concentration ( $\text{mg}/\text{m}^3$ ) in the eastern Mediterranean, Black Sea and Caspian Sea regions. Data after NASA/GSFC

being among the most troubled waters of the world ocean (IOC, 1993). Land use / cover and subsequent hydrological changes in the adjacent lands lead to desertification and scarcity of water (Moreno and Oechel, 1995; Jeftic, 1992, 1996; Glanz and Zonn, 1997), amplified by cultural and socio-economic contrasts in the region.

At present the causal relationships explaining the evolution of the ocean-atmosphere system projected onto the region are not well established. Often the changes are mediated through sub-basin and meso-scale processes, and are therefore difficult to be identified.

## 2. Large-Scale Controls

The region is one of the foremost areas of the world where interannual and long term climatic variability is predominant. In the Mediterranean region such variability is well known (e.g. Garrett *et al.*, 1992; Robinson *et al.*, 1993; Malanotte-Rizzoli and Robinson 1994; Jeftic *et al.*, 1992, 1996). The Mediterranean variability appears correlated with global teleconnection patterns, and coupled with the Indian Monsoon system and El Nino / Southern Oscillation (ENSO) (Ward, 1996). For example, good correlation has been found between heavy rain and snow in Israel during the last 100 years and ENSO (Alpert and Reisin, 1986). Similarly, global versus regional climate interaction has been emphasized in the case of the Caspian Sea (Rodionov, 1994) as well as in the Black Sea (Polonsky *et al.*, 1997).

Weather in Europe, extending well into Eurasia, is to a large extent determined by conditions in the North Atlantic, and in particular the North Atlantic Oscillation (NAO) quantified by the normalized anomaly of the sea level pressure difference between the Azores and Iceland (Hurrell, 1995). Severe weather in Europe occurs when the NAO index is positive. The NAO accounts for about one third of the hemispheric interannual variance, and accounts for surface temperature changes as well as evaporation-precipitation anomalies in the European and the Eastern Mediterranean regions (Hurrell, 1995, 1996; Marshall, 1997). It has been linked to sea level changes in the Caspian Sea (Rodionov, 1994), to Danube river runoff directly influencing Black Sea hydrology (Polonsky *et al.*, 1997), as well as to surface winter temperatures, precipitation and river runoff in Turkey (Cullen, 1998).

Large scale control is well expressed in long range atmospheric transport patterns, suggested by the simultaneous occurrence (Li *et al.*, 1996; Andreae, 1996) and parallel dependence on the interannual NAO patterns (Moulin *et al.*, 1997) of the transport of aerosol dust from the Sahara desert into the Mediterranean and tropical Atlantic regions. An exceptional case studied during the first half of April 1994 (Özsoy *et al.*, 1998; Kubilay *et al.*, 1998), has illustrated the role of large scale controls. Atmospheric blocking in the Atlantic had triggered upper air jet interactions and meridional circulations on a hemispherical scale (Figure 2a-c). These interactions resulted in large scale subsidence and cyclogenesis resulting in an anomalous pattern of dust simultaneously transported towards the subtropical Atlantic and the eastern Mediterranean regions, leading to maximum dust concentrations in 30 years of measurements in Bermuda and similarly high values in Erdemli (Özsoy, *et al.*, 1998). It is most interesting that the average sea level pressure was characterized with the typical dipole pattern of the NAO (Fig 2d), with a corresponding high index value of  $\sim 4$ , suggesting significant




Figure 2. Northern hemisphere upper atmospheric circulation showing the evolution of the polar and subtropical jets. 250hPa wind speed (shading) and direction (vectors) based on two day averages for (a) March 31 - April 1, (c) April 4 - 5, 1994, (c) April 12 - 13, 1994 and (d) the monthly average sea level pressure for the period March 15 - April 15, 1994. The source for the data is NOAA Climate Diagnostic Center (plotting page web address <http://www.cdc.noaa.gov/HistData/>)

links between high index NAO circulation and the anomalous dust event. In addition to the links with NAO, the Atlantic dust flux also appears well correlated with the African drought and the ENSO (Prospero and Nees, 1986).

### 3. Similarities in Regional Cooling Patterns

There is a significant degree of synchronism displayed between the Levantine, Black and Caspian Seas, in terms of the air and sea surface temperatures displayed in Figure 3. This is because of the proximity of the three regions, but also a result of the possible large scale controls discussed above, and by Özsoy and Ünlüata (1997). In the same Figure, comparisons are also made with the NAO and SO indices and with solar transmission (an indicator of volcanic dust in the atmosphere). Some cold years appear




Figure 3. Time series of average air (ECMWF/ERA re-analyses, 6 hr forecast temperature at 10m height) and sea surface temperatures (ESA ERS1/ATSR derived monthly averages) for the Levantine, Black and Caspian Seas, Solar transmission at Mauna Loa, and the climatic indices NAOI (seasonal averages) and SOI, for the last two decades.

linked with negative values of the SO Index in Figure 3 (e.g. 1982-83, 1986-87, the 1990's), often cited as ENSO years (e.g. Meyers and O'Brien, 1995). Similarly, some years (1983, 1986-87, 1989-90, 1992-93) are characterized by high NAO indices.

Relatively cooler winters are detected in the years 1982-83, 1985, 1987, 1989, 1991 and 1992-93 in Figure 3. Some of the cold years correspond to well known cases of convection and deep water formation in the regional seas (e.g. from recent data in 1987: dense water intrusion into the Marmara Sea from the Aegean, Beşiktepe *et al.*, 1993; deep water formation in the Rhodes Gyre region, Gertman *et al.*, 1990; main pycnocline erosion in the Black Sea, unpublished data; in 1989: extensive LIW formation in




Figure 4. Average lower tropospheric temperature anomalies for (a) the globe, (b) the Rhodes Gyre region of the Levantine Sea ( $33^{\circ}$ - $37^{\circ}$ N and  $26^{\circ}$ - $32^{\circ}$ E), (c) southwest Black Sea ( $41^{\circ}$ - $43^{\circ}$ N and  $27^{\circ}$ - $31^{\circ}$ E), (d) northern Caspian Sea ( $45^{\circ}$ - $47^{\circ}$ N and  $48^{\circ}$ - $53^{\circ}$ E). The measurements were obtained from the Microwave Sounding Unit (MSU) for the 0-5.6 km layer of the troposphere (grid resolution  $2.5^{\circ}$ ) and the anomalies calculated over the base period of 1979-1995 are produced by the Global Climate Perspectives System (GCPS) at the <http://www.ncdc.noaa.gov/onlineprod/prod.html> web address.

northern Levantine Sea (Özsoy *et al.*, 1993); in 1992 and 1993: Black Sea Cold Intermediate Water (CIW) formation and pycnocline erosion, Ivanov *et al.*, 1997a,b; simultaneous intermediate and deep water formation in the Rhodes Gyre region of the Levantine Sea; Sur *et al.*, 1992, Özsoy *et al.*, 1993). There are also some surprises: the year 1987 is one of the coolest years in all three seas, but there is no corresponding decrease of air temperature in the Caspian Sea. Secondly, while the air temperature displays various anomalous years in the Caspian Sea, the sea surface temperature does not respond to it very effectively, except the year 1987, unlike the pattern observed for the other two seas.

Cooling in the lower troposphere (Figure 4a) occurred globally in 1992-1993 (Spencer and Christy, 1992), following other cooling periods of 1982, 1985-86, 1989 in the last two decades. Significant drops of Temperature

drops of  $1$ - $2^{\circ}\text{C}$  occurred in the entire region extending from the Rhodes Gyre of the Levantine Sea to the southern Black Sea and the northern Caspian Sea (Figures 4b-d). An event of global climatic significance (Fiocco *et al.*, 1996) the eruption of Mount Pinatubo volcano in June 1991, resulting in stratospheric warmings (Angell, 1993), decreased solar energy inputs (Dutton and Christy, 1992; Dutton, 1994) and anomalous temperatures (Halpert *et al.*, 1993; Boden *et al.*, 1994) in the entire northern hemisphere in 1992-93. Anomalous cold temperatures appeared in the Middle East in very similar spatial patterns during the winters of 1992 and 1993, covering the Black Sea, eastern Mediterranean, and African regions in both years (Özsoy and Ünlüata, 1997). In Turkey, the winter of 1992 was the coldest in the last 60 years (Türkeş *et al.*, 1995), and in Israel, it was the coldest in the last 46 years (Genin *et al.*, 1996).

#### 4. Surface Fluxes

To study the effects of climate variability in the three seas, the loss terms of the surface fluxes are computed from uniform quality, decadal atmospheric re-analysis data obtained from the ECMWF at  $1^{\circ}$  resolution (mean sea level pressure, 10 m wind, 2 m atmospheric and dew point temperatures, and cloudiness, produced every 6 hr intervals from 6 hr global forecasts), and monthly average sea surface temperature based on ERS1/ATSR satellite data for the period 1981-1994. The air-sea fluxes are calculated by a method of iteratively reconstructing the atmospheric variables at 10 m height from ECMWF supplied fields based on Monin-Obukhov turbulent boundary layer theory, then using bulk formulae to compute the wind-stress, moisture flux, sensible ( $q_s$ ) and latent ( $q_l$ ) heat fluxes (Launiainen and Vihma, 1990; Vihma, 1995) as well as the longwave radiation ( $q_b$ ) fluxes (Bignami *et al.*, 1995). The values are then averaged over the sea domain and a month to produce the values in Figure 5, where the standard deviations are also marked.

The comparison of fluxes in the three different seas shows coincidence between the active periods of Black Sea and Levantine Sea, and a lesser degree of synchronism between them and the Caspian Sea. On the other hand, the Black Sea and Caspian Sea momentum fluxes are larger, more seasonal and more variable than the Levantine Sea, where the larger events come in interannual pulses. The sensible heat flux increases from the Levantine Sea to the Black Sea, reaching a maximum in the Caspian Sea.




Figure 5. Wind stress and heat flux components for the Levantine, Black and Caspian Seas computed from the ECMWF re-analysis, 6 hourly forecast fields. The values are averaged over each basin and over one month periods. The error bars denote one  $\sigma$  standard deviation.

## 5. Changes in the Eastern Mediterranean

### 5.1. THERMOHALINE CIRCULATION

The mean residence time varies considerably from  $\sim 100$  years for the Mediterranean and shorter for the Caspian, to  $\sim 2000$  for the Black Sea. The




Figure 6. Areas chosen for deep water analyses in the western Levantine, eastern Ionian and Cretan Seas

Mediterranean 'conveyor belt' (*i.e.* the 3-D thermohaline circulation, partly connected to the North Atlantic through the Gibraltar Strait), could have global impact, by regulating water masses and overturning in the North Atlantic (Reid *et al.*, 1994; Hecht *et al.*, 1996), with a potential, yet speculative contribution (Johnson, 1997a,b) to the triggering of paleoclimatic transitions (Broecker *et al.*, 1985).

The conceptual schemes of Mediterranean thermohaline circulations (*e.g.* Robinson and Golnaraghi, 1994) have been drastically changed by the discovery of deep water formation (a) through dense water outflow from the Aegean Sea, and (b) simultaneously with LIW in the Rhodes Gyre core. It is now evident that the 'conveyor belt' circulation in the entire Mediterranean is undergoing changes. Increases in the deep water temperature and nutrients (Bethoux *et al.*, 1989; Bethoux, 1989, 1993) appear coupled to the annual deep convection patterns (Gascard, 1991; Madec and Crepon, 1991; Send *et al.*, 1996) in the western Mediterranean.

The real surprise has recently come from the eastern Mediterranean, where deep water was found to form in the center of the permanent Rhodes Gyre, simultaneously with LIW on its periphery (Gertman *et al.*, 1990; Sur *et al.*, 1992; Özsoy *et al.*, 1993) with a recurrence interval of a few years depending on cooling. Furthermore, a climatically induced switching in the closed cell 'conveyor belt' is now evident. The classical scheme of deep water renewal by dense water (fresh, cold) outflow from the Adriatic Sea (Roether and Schlitzer, 1991; Schlitzer *et al.*, 1991; Roether *et al.*, 1994), have been replaced by the dense water (salty, warm) outflow from the Aegean Sea (Roether *et al.*, 1996), starting in the early 1990's. The event has been detected for first time since the beginning of oceanographic observations,




Figure 7. Average salinity, potential temperature, and log number of observations in the depth interval of 2000-4000m in the western Levantine, eastern Ionian Sea areas. The error bars denote standard deviation. The averages are obtained from individual data sets contained within the combined MODB / POEM data and grouped into 1 year intervals falling within the specified depth range.

although intermediate depth ( $\sim 1,000$ - $1500$ m) Aegean intrusions of lesser magnitude were well known (Roether and Schlitzer, 1991).

The changes in deep water first became evident when an unusually warm, saline water mass below  $1000$ m was detected south of the Island of Crete during summer 1993 (Heike *et al.*, 1994; A. Yilmaz, pers. comm.). Anomalous heat fluxes were detected in the deep sediments of Ionian Sea deep brine lakes in 1993-94, capped by a double diffusive interface in contact with the new deep water (Della Vedova *et al.*, 1995), suggesting a fresh transient event. Although an exact date for the Aegean dense water outflow can not be established, observations suggest a strong pulse in 1992-93, superposed on a background trend starting in the late 1980's.




Figure 8. Average salinity, potential temperature,  $\sigma_\theta$  density and log number of observations in the depth interval of 1000-2000m in the Cretan Sea. The error bars denote standard deviation. The averages are obtained from individual data sets contained within the combined MODB / POEM data and grouped into 1 year intervals falling within the specified depth range.

An analysis of the MODB historical data base (Brasseur *et al.*, 1996) combined with the last 10 years of Physical Oceanography of the Eastern Mediterranean (POEM) data was made by Özsoy and Latif (1996) to address the effect of Aegean outflow in the observed changes in deep water. For this purpose, the three regions adjacent to the Aegean - Eastern Mediterranean junction (Figure 6) were selected.

The average properties in the depth range of 2000-4000m, with the 95 % confidence limits are shown in Figure 7 the western Levantine and eastern Ionian boxes (Figure 6), together with the number of observations in each annual cluster. The vertical resolution, accuracy and coverage of the measurements increased in the second half of 1980's (POEM program). If we disregard the large salinity changes associated with measurement drift in the 1960's and 1970's, and trust the relatively more accurate temperature measurements, an increase in deep water temperature (and possibly salinity) is evident in both regions in the late 1980's, topped by a rapid increase during 1992-95.

In the Cretan Sea deep water (maximum depth  $\sim 2000$ m), both tem-

perature and salinity have fluctuated at least twice between minimum and maximum values in the last 60 years, starting a steady increase after the mid 1980's (Figure 8). The salinity reached its peak in 1995, while the temperature first increased until 1991 and dropped sharply in 1991-95 to yield a rapid and steady increase in density during the 1980's and 1990's, which is quite different than the relatively constant density values of the previous three decades. Similar conclusions were reached by Theocharis *et al.*, (1996) with regard to Cretan Sea.

## 5.2. NEAR-SURFACE CIRCULATION

Upper ocean variability has been most readily detected in the circulation of the main thermocline and in the physical properties of the intermediate and surface waters both in the western Mediterranean (Crepon *et al.*, 1989; Barnier *et al.*, 1989) and the eastern Mediterranean (Hecht, 1992; Sur, *et al.*, 1992; Özsoy *et al.*, 1993).

In the eastern Mediterranean, abrupt changes in circulation and water masses has been recognized (Hecht, 1992) as extraordinary, multi-annually recurrent events. The multiple bifurcating central jet flow joining the Asia Minor jet cyclonically east and west of Cyprus and anticyclonically along the eastern coast of Levant in 1985-86 has abruptly changed to a better organized, cyclonic flow around Cyprus in the 1988-1990 period, coincident with the Rhodes Gyre deep convection in 1987, the massive penetration of low salinity modified Atlantic water into the northern Levantine, and the disappearance of a coherent anticyclone in the Gulf of Antalya the same year and its re-appearance in 1990 (Özsoy *et al.*, 1991, 1993). The surface salinity steadily increased from 39.1-39.3 in 1985-86 up to 39.5 in 1989 and 39.7 in 1990. During the same period changes occurred in the pattern of formation and maintenance of LIW. The LIW trapped in the Antalya anticyclonic eddy in 1985-86 disappeared together with the eddy itself in 1987. LIW with increasingly higher salinities was observed in the winter periods of the following years, reaching from ~39.0 in 1989 to ~39.2 in 1989-90 (Özsoy *et al.*, 1993). An increasing trend was evident in the Shikmona Gyre core salinity, temperature and density during the 1988-1994 period, with abrupt changes in the winters of 1990 and 1992 (Brenner, 1996). Salinity increases in the entire region, together with circulation changes leading to the entry of salt water into the Aegean, and blocking of LIW to spread westward by multi-centered anticyclonic region in the Southern Levantine, have been shown by Malanotte-Rizzoli *et al.*, (1998) for the POEM survey of October 1991. The changes imply a salt redistribution pattern favoring the creation of Aegean dense water outflow.

## 6. Hydrological Cycles and Sea Level

The sea-level, besides being a good indicator of climatic fluctuations, is a sensitive measure of hydrometeorological driving factors in enclosed and semi-enclosed seas. In the Black Sea, sea-level responds non-isostatically to atmospheric pressure and the total water budget, which are both highly variable on interannual and seasonal time scales (Sur *et al.*, 1994), strictly controlled by the variable inputs of large rivers and the dynamical constraints imposed by the Turkish Straits (Özsoy *et al.*, 1998a). Simple models used to understand the time dependent response has had limited success to produce and explain sea-level variations of large amplitude in this non-tidal Sea, unless special effects of wind-setup on the hydraulics of the Bosphorus are included (Ducet and Le Traon, 1998).

In the Caspian Sea, sea-level changes depend on the regional hydrometeorological regime linked to global climate (Radionov, 1994). The sea level changes with climatic and anthropogenic components are of great economic and environmental importance for the surrounding countries. Interestingly, the sea-level change influences the residence time of the deep waters, and therefore has a direct bearing on the health of the Caspian Sea. Abrupt changes in sea-level have occurred twice since 1830's, as well as earlier in history, as the fate of Khazars occupying in its shores in the 10th century stand witness. The sea level has first dropped from a -25 m in 1930's down to -29 m by 1978, and has risen to the present -27 m after 1977. Hydrogen sulfide has been detected in deep waters prior to 1930 when sea-level was high, as a result of insufficient ventilation, limited by the decreased volume of dense water formed on the ice-covered, shallow northern Caspian shelf (average depth ~ 2m) under the influence of the large, variable inputs of the Volga river (Kosarev and Yablonskaya, 1994).

## 7. Marine Ecosystems

The marine ecosystems of interior seas are especially vulnerable, as significant changes in nutrient supplies are taking place in their confined waters. In the Black Sea, and the Caspian Sea, supplied by large rivers, eutrophication is leading to losses of habitats, species diversity, and consequent economical value (Zaitsev and Mamayev, 1997; Kosarev and Yablonskaya, 1994). The eastern Mediterranean system may be undergoing change related to the Nile river (*e.g.* Turley, 1997). Ecosystem disasters in even smaller, confined waters of the the Aral Sea, Kara Boğaz Göl, and Azov Sea, are better recognized. The introduction of extraneous species represent anthropogenic effects, either by filling a niche as in the case of the Black Sea, or through Lessepsian migration from the Red Sea, in the case of the eastern Mediterranean (Por, 1978; Galil, 1993; Gücü *et al.*, 1994;




Figure 9. Model derived tropospheric dust load ( $\text{g/m}^2$ ) at 18:00 on April 6, 1994.

Kıdeys and Gücü, 1995). On the other hand, the natural variabilities are frequently of the same magnitude as anthropogenic effects, which makes diagnosis difficult.

The basic ingredients and biological machinery of the marine ecosystems are often not sufficiently resolved (*e.g.* the sub-oxic zone in the Black Sea, Oğuz, *et al.*, *this volume*; or the roles of P-limitation and cyanobacteria in the eastern Mediterranean, Zohary and Robarts, 1996; Li *et al.*, 1993). In addition to these biochemical factors, dynamical features are of first order importance for the ecosystem. In the Black Sea, upwelling, coastal flows and river supply play important roles (Sur *et al.*, 1994, 1996; Özsoy and Ünlüata, 1997). In the eastern Mediterranean, where the surface nutrients are low, nutrient upwelling at fronts (Özsoy *et al.*, 1993), and nutrient supply in the river mouths and coasts appear significant. Ecosystem changes associated with an uplift of the nutricline following the Aegean dense water outflow are suspected in the eastern Mediterranean (Roether *et al.*, 1996), and are perhaps associated with changes in deep zooplankton (Weikert, 1996). The strong cooling and mixing in 1992-93 produced a massive algal bloom near the Rhodes Gyre (Yılmaz *et al.*, 1996; Ediger and Yılmaz, 1996), and in Eilat, where it led to the destruction of coral reefs (Genin *et al.*, 1996). These events coincided with the enhanced pycnocline erosion (Ivanov *et al.*, 1997a,b), followed by a massive bloom (Vladimirov *et al.*, 1997) in the following summer of 1992 in the Black Sea. The bloom in the Black Sea apparently produced atmospheric non-sea-salt sulfate aerosols of marine biological origin swept southwards and detected in the Erdemli

measurements (Özsoy *et al.*, 1998c).

## 8. Aerosol Dust

Aerosol desert dust is important in the climate system, primarily because of its effects on heat budgets (Charlson and Heintzenberg, 1995), especially in semi-enclosed seas (Gilman and Garrett, 1994), on the heterogeneous chemistry of the tropospheric and greenhouse gases (Dentener *et al.*, 1996), as well as on the biogeochemical cycles in the marine environment (Duce *et al.*, 1991).

Dust transport is an important climatic process which is particularly active in the Mediterranean area (Guerzoni and Chester, 1996), with an increasing incidence in recent years associated with drought and desertification. The role of large scale controls in establishing conditions for transport, such as in the spectacular April 1994 case, are outlined in Section 2. The dust load for this unique case of simultaneous transport (Andrae, 1996; Moulin *et al.*, 1997) in a shallow layer across the Atlantic and within a rapidly developing cyclone towards the eastern Mediterranean and Black Sea regions, resulting in anomalously high concentrations in Barbados and Erdemli (Li *et al.*, 1996, Özsoy *et al.*, 1998b; Kubilay *et al.*, 1998) is shown in Figure 9.

Sahara dust is known to be the main source contributing to the formation of the fertile 'red soils' of the eastern Mediterranean. It is hypothesized, based on satellite observations and concurrent dust measurements (unpublished material, IMS-METU) that the transport and deposition on sea surface of eroded dust from the Sahara and Arabian deserts has an impact on short term, episodic phytoplankton blooms in the eastern Mediterranean. There is often a striking coincidence of 'high reflectivity' from the sea surface typically associated with *E. huxleyi* blooms, and the incursions of dust, supported by the coincidence of the fertilization events with high fluxes of carbonates in sediment trap observations. However, the working hypothesis has to be better defined and tested with detailed observations.

## 9. ACKNOWLEDGEMENTS

This study benefited from a number of activities, most notably the POEM and POEM-BC, international cooperative programs, the NATO Sfs Projects TU-FISHERIES, TU-BLACK SEA and TU-REMOSSENS projects, as well as other work carried out by the IMS-METU. Special thanks are for the Turkish General Directorate of Meteorology for providing access to the ECMWF data on a collaborative basis.

42. Johnson, R. G. (1997a). Ice Age Initiation by an Ocean-Atmosphere Circulation Change in the Labrador Sea, *Earth Planet. Sci. Lett.*, **148**, 367-379.
43. Johnson, R. G. (1997b). Climate Control Requires a Dam at the Strait of Gibraltar, *EOS*, July 8, 1997.
44. Kıdeys, A. E. and A. C. Gücü (1995). Rhophilema nomadica: a Poisonous Indo-Pacific Scyphomedusan New to the Mediterranean Coast of Turkey, *Israel J. of Zoology*, **41**(4): 615- 617.
45. Kosarev, A. N. and E. A. Yablonskaya (1994). *The Caspian Sea*, Backhuys Publishers, Hague, 259 pp.
46. Kubilay, N., E. Özsoy, S. Nickovic, I. Tegen and C. Saydam (1998). A Hemispheric Dust Storm Affecting the Atlantic and Mediterranean (April 1994): Analyses, Modelling, Ground-Based Measurements and Satellite Observations, (submitted).
47. Launiau, J. and T. Vihma (1990). Derivation of Turbulent Surface Fluxes - An Iterative Flux-Profile Method Allowing Arbitrary Observing Heights, *Environmental Software*, **5**(3), 113-124.
48. Li, W. K. W., T. Zohary, Y. Z. Yacobi and A. M. Wood (1993). Ultraphytoplankton in the Eastern Mediterranean Sea: Towards Deriving Phytoplankton Biomass from Flow Cytometric Measurements of Abundance, Fluorescence and Light Scatter, *Marine Ecology Progress Series*, **102**, 79-87.
49. Li, X., H. Maring, D. Savoie, K. Voss and J. M. Prospero (1996). Dominance of Mineral Dust in Aerosol Light Scattering in the North Atlantic Trade Winds, *Nature*, **380**, 416-419.
50. Malanotte-Rizzoli, P. and A. R. Robinson (editors) (1994). *Ocean Processes in Climate Dynamics: Global and Mediterranean Examples*, NATO ASI Series C: Mathematical and Physical Sciences, vol. 419, Kluwer Acad. Publ., Dordrecht, 437 pp.
51. Malanotte-Rizzoli, P., B. B. Manca, M. Ribera d'Alcala, A. Theocharis, S. Brenner, G. Budillon and E. Özsoy (1998). The Eastern Mediterranean in the 80s and in the 90s: The Big Transition in the Intermediate and Deep Circulations (submitted).
52. Madec, G. and M. Crepon (1991). Thermohaline-Driven Deep Water Formation in the Northwestern Mediterranean Sea, in: P. C. Chu and J. C. Gascard (editors) *Deep Convection and Deep Water Formation in the Oceans*, Elsevier, Amsterdam, pp. 241-265.
53. Marshall, J. and Y. Kushnir (1997). A 'white paper' on Atlantic Climate Variability, an unpublished working document.
54. Meyers, S. D. and J. J. O'Brien (1995). Pacific Ocean Influences Atmospheric Carbon Dioxide, *EOS*, **76**(52), 533-537.
55. Moreno, J. M. and Oechel, W., Eds. (1995). *Global Change and Mediterranean Ecosystems*, Springer Verlag, Ecological Studies, **117**, New York, N.Y.
56. Moulin, C., C. E. Lambert, F. Dulac and U. Dayan (1997). Control of atmospheric export of dust from North Africa by the North Atlantic oscillation. *Nature*, **387**, 691-694.
57. Özsoy, E. (1981). *On the Atmospheric Factors Affecting the Levantine Sea*, European Center for Medium Range Weather Forecasts, Reading, U.K., Technical Report No. 25, 30 pp.
58. Özsoy, E., A. Hecht, Ü. Ünlüata, S. Brenner, T. Oğuz, J. Bishop, M. A. Latif and Z. Rozentraub (1991). A Review of the Levantine Basin Circulation and its Variability during 1985-1988, *Dyn. Atmos. Oceans*, **15**, 421-456.
59. Özsoy, E., Hecht, A., Ünlüata, Ü., Brenner, S., Sur, H. İ., Bishop, J., Latif, M. A., Rozentraub, Z. and T. Oğuz (1993). A Synthesis of the Levantine Basin Circulation and Hydrography, 1985-1990, *Deep-Sea Res.*, **40**, 1075-1119.
60. Özsoy, E. and M. A. Latif (1996). Climate Variability in the Eastern Mediterranean and the Great Aegean Outflow Anomaly, International POEM-BC/MTP Symposium, Molitg les Bains, France, 1-2 July 1996, pp. 69-86.
61. Özsoy, E. and Ü. Ünlüata (1997). Oceanography of the Black Sea: A Review of Some Recent Results, *Earth Sci. Rev.*, **42**(4), 231-272.
62. Özsoy, E., Latif, M. A., Beşiktepe, S., Çetin, N., Gregg, N. Belokopytov, V., Goryachkin, Y. and V. Diaconu (1998a). The Bosphorus Strait: Exchange Fluxes, Currents and Sea-Level Changes, L. Ivanov and T. Oğuz, editors, *Proceedings of NATO TU-Black Sea Project - Ecosystem Modeling as a Management Tool for the Black Sea: Symposium on Scientific Results, Crimea, (Ukraine), June 15-19, 1997*.
63. Özsoy, E., N. Kubilay, S. Nickovic and C. Saydam (1998b). A Hemispheric Dust Storm in April 1994: Observations, Modelling and Analyses, XXIIIrd NATO/CCMS International Technical Meeting on Air Pollution, Varna, 28 Sep. - 2 Oct. 1998. (submitted for publication).
64. Özsoy, T., C. Saydam, N. Kubilay, O. B. Nalçacı and İ. Salihoğlu (1998c). Aerosol Nitrate and Non-Sea-Salt Sulfate Over the Eastern Mediterranean, XXIIIrd NATO/CCMS International Technical Meeting on Air Pollution, Varna, 28 Sep. - 2 Oct. 1998. (submitted for publication).
65. Polonsky, A., E. Voskresenskaya and V. Belokopytov, Variability of the Northwestern Black Sea Hydrography and River Discharges as Part of Global Ocean-Atmosphere Fluctuations, in: E. Özsoy and A. Mikaelyan (editors), *Sensitivity to Change: Black Sea, Baltic Sea and North Sea*, NATO ASI Series (Partnership Sub-series 2, Environment, 27), Kluwer Academic Publishers, Dordrecht, 536 pp., 1997.
66. Por, F. D. (1978). Lessepsian Migration - The influx of Red Sea Biota into the Mediterranean by way of the Suez Canal, Springer Verlag, 215 pp.
67. Reid, J. L. (1994). On the Total Geostrophic Circulation of the North Atlantic Ocean: Flow Patterns, Tracers and Transports, *Prog. Oceanogr.*, **33**, 1-92.
68. Reiter, E. R. *Handbook for Forecasters in the Mediterranean; Weather Phenomena of the Mediterranean Basin*; Part 1: General Description of the Meteorological Processes, Environmental Prediction Research Facility, Naval Postgraduate School, Monterey, California, Technical Paper No. 5-75, 344pp, 1979.
69. Robinson, A. R., Garrett, C. J., Malanotte-Rizzoli, P., Manabe, S., Philander, S. G., Pinardi, N., Roether, W., Schott, F. A., and J. Shukla (1993). Mediterranean and Global Ocean Climate Dynamics, *EOS*, **74**(44), November 2, 1993, 506-507.
70. Robinson, A. R. and M. Golnaraghi (1994). The Physical and Dynamical Oceanography of the Mediterranean Sea, in: Malanotte-Rizzoli P. and A. R. Robinson (editors), *Ocean Processes in Climate Dynamics: Global and Mediterranean Examples*, NATO ASI Series C: Mathematical and Physical Sciences, vol. 419, Kluwer Academic Publishers, Dordrecht, pp. 255-306.
71. Rodionov, S. N., *Global and Regional Climate Interaction: The Caspian Sea Experience*, Water Science and Technology Library, v. 11, Kluwer Academic Publishers, Dordrecht, 256 pp., 1994.
72. Roether, W. and R. Schlitzer (1991). Eastern Mediterranean Deep Water Renewal on the Basis of Chlorofluoromethane and Tritium Data, *Dyn. Atmos. Oceans*, **15**, 333-354, 1991.
73. Roether, W. V. M. Roussenov and R. Well (1994). A Tracer Study of the Thermohaline Circulation of the Eastern Mediterranean, in: P. Malanotte-Rizzoli and A. R. Robinson (editors), *Ocean Processes in Climate Dynamics: Global and Mediterranean Examples*, NATO ASI Series C: Mathematical and Physical Sciences, vol. 419, Kluwer Academic Publishers, Dordrecht, pp. 371-394.
74. Roether, W., Manca, B., Klein, B., Bregant, D., Georgopoulos D., Beitzel V., Kovacevic, V., and A. Luchetta (1996). Recent Changes in Eastern Mediterranean Deep Waters, *Science*, **271**, 333- 335.
75. Schlitzer, R., Roether, W., Oster, H., Junghans, H.-G., Hausmann, M., Johannsen H. and A. Michelato (1991). Chlorofluoromethane and Oxygen in the Eastern Mediterranean, *Deep-Sea Res.*, **38**, 1351-1551.
76. Send, U., J. Font and C. Mertens (1996). Recent Observation Indicates Convection's Role in Deep Water Circulation, *EOS*, **77**(7), February 13, 1996, 61-65.

42. Johnson, R. G. (1997a). Ice Age Initiation by an Ocean-Atmosphere Circulation Change in the Labrador Sea, *Earth Planet. Sci. Lett.*, **148**, 367-379.
43. Johnson, R. G. (1997b). Climate Control Requires a Dam at the Strait of Gibraltar, *EOS*, July 8, 1997.
44. Kıdeys, A. E. and A. C. Gücü (1995). Rhophilema nomadica: a Poisonous Indo-Pacific Scyphomedusan New to the Mediterranean Coast of Turkey, *Israel J. of Zoology*, **41**(4): 615- 617.
45. Kosarev, A. N. and E. A. Yablonskaya (1994). *The Caspian Sea*, Backhuys Publishers, Hague, 259 pp.
46. Kubilay, N., E. Özsoy, S. Nickovic, I. Tegen and C. Saydam (1998). A Hemispheric Dust Storm Affecting the Atlantic and Mediterranean (April 1994): Analyses, Modelling, Ground-Based Measurements and Satellite Observations, (submitted).
47. Launiau, J. and T. Vihma (1990). Derivation of Turbulent Surface Fluxes - An Iterative Flux-Profile Method Allowing Arbitrary Observing Heights, *Environmental Software*, **5**(3), 113-124.
48. Li, W. K. W., T. Zohary, Y. Z. Yacobi and A. M. Wood (1993). Ultraphytoplankton in the Eastern Mediterranean Sea: Towards Deriving Phytoplankton Biomass from Flow Cytometric Measurements of Abundance, Fluorescence and Light Scatter, *Marine Ecology Progress Series*, **102**, 79-87.
49. Li, X., H. Maring, D. Savoie, K. Voss and J. M. Prospero (1996). Dominance of Mineral Dust in Aerosol Light Scattering in the North Atlantic Trade Winds, *Nature*, **380**, 416-419.
50. Malanotte-Rizzoli, P. and A. R. Robinson (editors) (1994). *Ocean Processes in Climate Dynamics: Global and Mediterranean Examples*, NATO ASI Series C: Mathematical and Physical Sciences, vol. 419, Kluwer Acad. Publ., Dordrecht, 437 pp.
51. Malanotte-Rizzoli, P., B. B. Manca, M. Ribera d'Alcala, A. Theocharis, S. Brenner, G. Budillon and E. Özsoy (1998). The Eastern Mediterranean in the 80s and in the 90s: The Big Transition in the Intermediate and Deep Circulations (submitted).
52. Madec, G. and M. Crepon (1991). Thermohaline-Driven Deep Water Formation in the Northwestern Mediterranean Sea, in: P. C. Chu and J. C. Gascard (editors) *Deep Convection and Deep Water Formation in the Oceans*, Elsevier, Amsterdam, pp. 241-265.
53. Marshall, J. and Y. Kushnir (1997). A 'white paper' on Atlantic Climate Variability, an unpublished working document.
54. Meyers, S. D. and J. J. O'Brien (1995). Pacific Ocean Influences Atmospheric Carbon Dioxide, *EOS*, **76**(52), 533-537.
55. Moreno, J. M. and Oechel, W., Eds. (1995). *Global Change and Mediterranean Ecosystems*, Springer Verlag, Ecological Studies, **117**, New York, N.Y.
56. Moulin, C., C. E. Lambert, F. Dulac and U. Dayan (1997). Control of atmospheric export of dust from North Africa by the North Atlantic oscillation. *Nature*, **387**, 691-694.
57. Özsoy, E. (1981). *On the Atmospheric Factors Affecting the Levantine Sea*, European Center for Medium Range Weather Forecasts, Reading, U.K., Technical Report No. 25, 30 pp.
58. Özsoy, E., A. Hecht, Ü. Ünlüata, S. Brenner, T. Oğuz, J. Bishop, M. A. Latif and Z. Rozentraub (1991). A Review of the Levantine Basin Circulation and its Variability during 1985-1988, *Dyn. Atmos. Oceans*, **15**, 421-456.
59. Özsoy, E., Hecht, A., Ünlüata, Ü., Brenner, S., Sur, H. İ., Bishop, J., Latif, M. A., Rozentraub, Z. and T. Oğuz (1993). A Synthesis of the Levantine Basin Circulation and Hydrography, 1985-1990, *Deep-Sea Res.*, **40**, 1075-1119.
60. Özsoy, E. and M. A. Latif (1996). Climate Variability in the Eastern Mediterranean and the Great Aegean Outflow Anomaly, International POEM-BC/MTP Symposium, Molitg les Bains, France, 1-2 July 1996, pp. 69-86.
61. Özsoy, E. and Ü. Ünlüata (1997). Oceanography of the Black Sea: A Review of Some Recent Results, *Earth Sci. Rev.*, **42**(4), 231-272.
62. Özsoy, E., Latif, M. A., Beşiktepe, S., Çetin, N., Gregg, N. Belokopytov, V., Goryachkin, Y. and V. Diaconu (1998a). The Bosphorus Strait: Exchange Fluxes, Currents and Sea-Level Changes, L. Ivanov and T. Oğuz, editors, *Proceedings of NATO TU-Black Sea Project - Ecosystem Modeling as a Management Tool for the Black Sea: Symposium on Scientific Results, Crimea, (Ukraine), June 15-19, 1997*.
63. Özsoy, E., N. Kubilay, S. Nickovic and C. Saydam (1998b). A Hemispheric Dust Storm in April 1994: Observations, Modelling and Analyses, XXIIIrd NATO/CCMS International Technical Meeting on Air Pollution, Varna, 28 Sep. - 2 Oct. 1998. (submitted for publication).
64. Özsoy, T., C. Saydam, N. Kubilay, O. B. Nalçacı and İ. Salihoğlu (1998c). Aerosol Nitrate and Non-Sea-Salt Sulfate Over the Eastern Mediterranean, XXIIIrd NATO/CCMS International Technical Meeting on Air Pollution, Varna, 28 Sep. - 2 Oct. 1998. (submitted for publication).
65. Polonsky, A., E. Voskresenskaya and V. Belokopytov, Variability of the Northwestern Black Sea Hydrography and River Discharges as Part of Global Ocean-Atmosphere Fluctuations, in: E. Özsoy and A. Mikaelyan (editors), *Sensitivity to Change: Black Sea, Baltic Sea and North Sea*, NATO ASI Series (Partnership Sub-series 2, Environment, 27), Kluwer Academic Publishers, Dordrecht, 536 pp., 1997.
66. Por, F. D. (1978). Lessepsian Migration - The influx of Red Sea Biota into the Mediterranean by way of the Suez Canal, Springer Verlag, 215 pp.
67. Reid, J. L. (1994). On the Total Geostrophic Circulation of the North Atlantic Ocean: Flow Patterns, Tracers and Transports, *Prog. Oceanogr.*, **33**, 1-92.
68. Reiter, E. R. *Handbook for Forecasters in the Mediterranean; Weather Phenomena of the Mediterranean Basin*; Part 1: General Description of the Meteorological Processes, Environmental Prediction Research Facility, Naval Postgraduate School, Monterey, California, Technical Paper No. 5-75, 344pp, 1979.
69. Robinson, A. R., Garrett, C. J., Malanotte-Rizzoli, P., Manabe, S., Philander, S. G., Pinardi, N., Roether, W., Schott, F. A., and J. Shukla (1993). Mediterranean and Global Ocean Climate Dynamics, *EOS*, **74**(44), November 2, 1993, 506-507.
70. Robinson, A. R. and M. Golnaraghi (1994). The Physical and Dynamical Oceanography of the Mediterranean Sea, in: Malanotte-Rizzoli P. and A. R. Robinson (editors), *Ocean Processes in Climate Dynamics: Global and Mediterranean Examples*, NATO ASI Series C: Mathematical and Physical Sciences, vol. 419, Kluwer Academic Publishers, Dordrecht, pp. 255-306.
71. Rodionov, S. N., *Global and Regional Climate Interaction: The Caspian Sea Experience*, Water Science and Technology Library, v. 11, Kluwer Academic Publishers, Dordrecht, 256 pp., 1994.
72. Roether, W. and R. Schlitzer (1991). Eastern Mediterranean Deep Water Renewal on the Basis of Chlorofluoromethane and Tritium Data, *Dyn. Atmos. Oceans*, **15**, 333-354, 1991.
73. Roether, W. V. M. Roussenov and R. Well (1994). A Tracer Study of the Thermohaline Circulation of the Eastern Mediterranean, in: P. Malanotte-Rizzoli and A. R. Robinson (editors), *Ocean Processes in Climate Dynamics: Global and Mediterranean Examples*, NATO ASI Series C: Mathematical and Physical Sciences, vol. 419, Kluwer Academic Publishers, Dordrecht, pp. 371-394.
74. Roether, W., Manca, B., Klein, B., Bregant, D., Georgopoulos D., Beitzel V., Kovacevic, V., and A. Luchetta (1996). Recent Changes in Eastern Mediterranean Deep Waters, *Science*, **271**, 333- 335.
75. Schlitzer, R., Roether, W., Oster, H., Junghans, H.-G., Hausmann, M., Johannsen H. and A. Michelato (1991). Chlorofluoromethane and Oxygen in the Eastern Mediterranean, *Deep-Sea Res.*, **38**, 1351-1551.
76. Send, U., J. Font and C. Mertens (1996). Recent Observation Indicates Convection's Role in Deep Water Circulation, *EOS*, **77**(7), February 13, 1996, 61-65.

77. Spencer, R. W. and J. R. Christy (1992). Precision and Radiosonde Validation of Satellite Gridpoint Temperature Anomalies. Part II: A Tropospheric Retrieval and Trends During 1979-1990, *J. Climate*, 5, 858-866.
78. Sur, H. I., Özsoy, E., and Ü. Ünlüata (1992). Simultaneous Deep and Intermediate Depth Convection in the Northern Levantine Sea, Winter 1992, *Oceanol. Acta*, 16, 33-43.
79. Sur, H. I., Özsoy, E. and Ü. Ünlüata, (1994). Boundary Current Instabilities, Upwelling, Shelf Mixing and Eutrophication Processes In The Black Sea, *Prog. Oceanog.*, 33, 249-302.
- Sur, H. I., Y. P. Ilyin, E. Özsoy and Ü. Ünlüata, (1996). The Impacts of Continental Shelf / Deep Water Interactions in the Black Sea, *J. Mar. Sys.*, 7, 293-320.
80. Theocharis, A., H. Kontoyannis, S. Kioroglou and V. Papadopoulos (1996), International POEM-BC/MTP Symposium, Molitg les Bains, France, 1-2 July 1996, pp. 58-60.
81. Türkeş, M., Sümer, U., and G. Kılıç, 1995. Variations and Trends in Annual Mean Air Temperatures in Turkey With Respect To Climatic Variability, *International Journal of Climatology*, 15, 557-569.
82. Turley, C. M. (1997). The Changing Mediterranean Sea - A Sensitive Ecosystem ?, Abstracts Volume, International Conference on the Progress in Oceanography of the Mediterranean Sea, EC-MAST Programme, Rome, November 17-19, 1997.
83. Vihma, T. (1995). Atmosphere-Surface Interactions over Polar Oceans and Heterogeneous Surfaces, *Finnish Marine Res.* 264, 3-41.
84. Vladimirov, V. L., V. I. Mankovsky, M. V. Solov'ev and A. V. Mishonov (1997). Seasonal and Long-term Variability of the Black Sea Optical Parameters, in: E. Özsoy and A. Mikaelyan (editors), *Sensitivity to Change: Black Sea, Baltic Sea and North Sea*, NATO ASI Series (Partnership Sub-series 2, Environment, 27), Kluwer Academic Publishers, Dordrecht, 536 pp., 1997.
85. Wallace, J. M. and M. L. Blackmon, Observations of Low-frequency Atmospheric Variability, in: B. J. Hoskins, and R. P. Pearce, *Large-Scale Dynamical Processes in the Atmosphere*, Academic Press, 397 pp., 1983.
86. Ward, N. (1995). Local and Remote Climate Variability Associated with Mediterranean Sea-Surface Temperature Anomalies, European Research Conference on Mediterranean Forecasting, La Londe les Maures, France, 21-26 October 1995.
87. Weikert, H. (1996). Changes in Levantine Deep Sea Zooplankton, International POEM-BC/MTP Symposium, Molitg les Bains, France, 1-2 July 1996, pp. 99-101.
88. Yilmaz A., D. Ediger, Ç. Polat, S. Tuğrul, and İ. Salihoğlu (1996). The Effect of Cold and Warm Core Eddies on the Distribution and Stoichiometry of Dissolved Nutrients and Enhancement of Primary Production and Changes in Elemental Composition of Phytoplankton in the Eastern Mediterranean, International POEM-BC/MTP Symposium, Molitg les Bains, France, 1-2 July 1996, pp. 122-127.
89. Zaitsev, Yu. and V. Mamaev (1997). Marine Biological Diversity in the Black Sea, A Study of Change and Decline, Black Sea Environmental Series, Black Sea Environmental Series, Vol. 3, GEF Black Sea Environmental Programme, United Nations Publications, New York, 208 pp.
90. Zohary, T. and R. Robarts (1996). Experimental Study of Microbial P-limitation in the Eastern Mediterranean, International POEM-BC/MTP Symposium, Molitg les Bains, France, 1-2 July 1996, p. 128.

## SOME ASPECTS OF THE DARDANELLES

### 1. Introduction

This paper focuses on the physical characteristics of the brackish waters of the Aegean Sea. The Aegean Sea is a different physical system at the edge of the transformation of the Aegean Sea coastal system at the edge of the irregular coastline. The most complicated system is divided into the Aegean Sea coastal system, Chios Basin, and these parts of the system between the islands. The characteristics of the Aegean Sea coastal system are as follows:

The hydrological characteristics of the Mediterranean. The precipitation over the Aegean Sea is high throughout the year, the rate of evaporation is low. Brackish water flows form a well-organized density of Black Sea water flows comes from the Aegean Sea water rises above the Aegean Sea water. The turbulent mixing hampers the generation of salt water.