

DEVELOPMENT AND APPLICATIONS OF THE COASTAL QUASIGEOSTROPHIC  
OCEAN MODEL TO THE CASE OF MULTIPLE COASTS AND  
MULTIPLY CONNECTED DOMAINS

Emin Ozsoy<sup>\*</sup>, Carlos Lozano<sup>†</sup>, Ralph Milliff<sup>‡</sup> and Allan Robinson<sup>†</sup>

<sup>\*</sup>Institute of Marine Sciences, Middle East Technical  
University, P.K. 28, Erdemli, Icel, 33731 Turkey

<sup>†</sup>Harvard University, Division of Applied Sciences, Cambridge,  
MA 02138 USA

<sup>‡</sup>National Center for Atmospheric Research,  
Oceanography Section,  
Boulder, CO 80307 USA

The Harvard quasigeostrophic ocean model has been generalized to be applicable to semi-enclosed or closed ocean domains bounded by multiple segments of coasts and/or multiply connected geometry. Inviscid coastal boundary conditions and consistency constraints are required and the Greens function methods are utilized in the solution algorithm. The performance of the model is guaranteed by comparison with analytical solutions of Flierl (1977) and cases derived from them. The methodology is applied to the quasigeostrophic dynamical modeling of the Levantine Basin. The robustness of the method is illustrated by the stable preliminary predictions and sensitivity runs.