



## The role of atmospheric deposition in the biogeochemistry of the Mediterranean Sea<sup>☆</sup>

Stefano Guerzoni <sup>a,\*</sup>, Roy Chester <sup>b</sup>, François Dulac <sup>c</sup>,  
Barak Herut <sup>d</sup>, Marie-Dominique Loÿe-Pilot <sup>e</sup>,  
Chris Measures <sup>f</sup>, Christophe Migon <sup>g</sup>, Emanuela Molinaroli <sup>h</sup>,  
Cyril Moulin <sup>c</sup>, Paolo Rossini <sup>a</sup>, Cemal Saydam <sup>i</sup>,  
Alexandre Soudine <sup>j</sup>, Patrizia Ziveri <sup>k</sup>

<sup>a</sup> *Istituto di Geologia Marina, CNR/via Gobetti, 101/40129 Bologna, Italy*

<sup>b</sup> *Oceanography Laboratory, University of Liverpool, P.O. Box 147 Liverpool, L69 3BX, UK*

<sup>c</sup> *CEA, MIXTE Lab., Ctr. Faibles Radioactivites, Saclay 709-CNRS, 91198 Gif-sur-Yvette, France*

<sup>d</sup> *Israel Oceanographic and Limnological Research, Tel Shikmona, P.O. Box 8030, 31080 Haifa, Israel*

<sup>e</sup> *Ecole Normale Supérieure, IBM-CNRS 386/1, Rue Maurice Arnoux, 792120 Montrouge, France*

<sup>f</sup> *Dept. of Oceanography, University of Hawaii at Mānoa, 1000 Pope Road, Honolulu, HI 96 822, USA*

<sup>g</sup> *C.N.R.S., Laboratoire de Physique et Chimie Marines, B.P. 8/06230 Villefranche-sur-mer, France*

<sup>h</sup> *Dip. Scienze Ambientali, Università di Venezia, Dorsoduro 2137/30123 Venezia, Italy*

<sup>i</sup> *Institute of Marine Science-METU, P.O. Box 28, 33731 Erdemli-İçel, Turkey*

<sup>j</sup> *WMO, Environment Division, Case postale no. 2300/CH-1211, Geneva 2, Switzerland*

<sup>k</sup> *Geomarine Center, Vrije Universiteit Amsterdam, 1081 HV, Amsterdam, Netherlands*

---

### Abstract

Estimates of atmospheric inputs to the Mediterranean and some coastal areas are reviewed, and uncertainties in these estimates considered. Both the magnitude and the mineralogical composition of atmospheric dust inputs indicate that eolian deposition is an important (50%) or even dominant (>80%) contribution to sediments in the offshore waters of the entire Mediterranean basin. Model data for trace metals and nutrients indicate that the atmosphere delivers more than half the lead and nitrogen, one-third of total phosphorus, and 10% of the zinc entering the entire basin. Measured data in sub-basins, such as the north-western Mediterranean and northern Adriatic indicate an even greater proportions of atmospheric versus riverine

---

<sup>☆</sup> This paper is an expansion of discussions held at the CIESM-CNR workshop in Bologna, 8–11 October 1997.

\* Corresponding author.

*E-mail address:* guerzoni@igm.bo.cnr.it (S. Guerzoni)

inputs. When dissolved fluxes are compared (the form most likely to impinge on surface water biogeochemical cycles), the atmosphere is found to be 5 to 50 times more important than rivers for dissolved zinc and 15 to 30 times more important for lead fluxes. Neglecting co-limitation by other nutrients, new production supported by atmospheric nitrogen deposition ranges from 2–4 g C m<sup>-2</sup> yr<sup>-1</sup>, whereas atmospheric phosphorus deposition appears to support less than 1 g C m<sup>-2</sup> yr<sup>-1</sup>. In spite of the apparently small contribution of atmospheric deposition to overall production in the basin it has been suggested that certain episodic phytoplankton blooms are triggered by atmospheric deposition of N, P or Fe. Future studies are needed to clarify the extent and causal links between these episodic blooms and atmospheric/oceanographic forcing functions. A scientific program aimed at elucidating the possible biogeochemical effects of Saharan outbreaks in the Mediterranean through direct sampling of the ocean and atmosphere before and after such events is therefore highly recommended. © 1999 Elsevier Science Ltd. All rights reserved.

## Contents

1. Introduction . . . . .	149
2. Magnitude and mineralogical composition of atmospheric mineral dust input . . .	150
2.1. Atmospheric mineral particle fluxes and mineralogy . . . . .	152
2.1.1. Mineralogy . . . . .	153
2.2. Sediment trap estimates . . . . .	155
2.3. Remote sensing and modelling . . . . .	156
3. Magnitude and fate of atmospheric trace metal inputs . . . . .	157
3.1. Trace metal concentrations in air . . . . .	157
3.2. Metal fluxes (total, dry, wet) . . . . .	158
3.3. Fate of trace metals in seawater . . . . .	160
3.4. Atmospheric versus riverine inputs . . . . .	163
3.4.1. Measurements . . . . .	163
3.4.2. Models . . . . .	164
3.5. Case studies . . . . .	167
3.5.1. The atmospheric 'signature' in seawater . . . . .	167
3.5.2. Pb in atmosphere and seawater . . . . .	169
4. Atmospheric input of nutrients: estimates and impacts . . . . .	169
4.1. Atmospheric inputs of nitrogen to the Mediterranean . . . . .	170
4.1.1. Atmospheric input of wet inorganic nitrogen . . . . .	170
4.1.2. Atmospheric input of dry inorganic nitrogen . . . . .	171
4.1.3. Total inorganic nitrogen deposition (Table 14) . . . . .	171
4.1.4. Organic nitrogen . . . . .	172
4.1.5. Atmospheric versus riverine inputs of nitrogen in the Mediterranean . . . .	173
4.1.6. Impact of atmospheric input of nitrogen on primary production . . . . .	173
4.2. Atmospheric input of phosphorus . . . . .	174
4.2.1. Saharan dust and phosphorus . . . . .	174
4.2.2. Impact of atmospheric phosphorus on primary production . . . . .	175
4.3. Conclusions and hypotheses . . . . .	176

5. Potential effects of atmospheric input on surface ocean biogeochemistry: preliminary speculations . . . . .	177
5.1. Meteosat and CZAS: a case of P-fertilisation? . . . . .	177
5.2. Wet deposition of dust: the role of Fe speciation in phytoplankton blooms . . .	178
6. Conclusions . . . . .	180
6.1. Future work . . . . .	181
Acknowledgements . . . . .	184
References . . . . .	184

---

## 1. Introduction

Over the past two decades it has become apparent that the atmosphere is not only a significant but in some cases the dominant pathway by which both natural and pollutant materials are transported from continents to both coastal and open seas. These substances include mineral dust, plant residues, heavy metals, nitrogen species from combustion processes and fertilisers, pesticides, and a wide range of synthetic organic compounds from industrial and domestic sources. Some of these substances, such as lead and some chlorinated hydrocarbons, when carried into oceans by the atmosphere, are potentially harmful to marine biological systems. Others, such as nitrogen species, phosphorus and iron, are nutrients and may enhance marine productivity. For some substances (e.g. aluminium, cadmium, lead, and some rare earth elements), atmospheric flux has an important impact on their oceanic chemical cycles. Atmospheric inputs may thus exert many different effects on the marine environment, and it is necessary to understand the magnitude and geographical distribution of the atmospheric fluxes of these materials to both coastal and open seas. Indeed, there have recently been a number of intensive studies of atmospheric inputs to coastal waters in North America and Europe, with several studies also devoted to investigating these processes in open oceans.

The most complete review of this subject, updated to the early 1990s, was a report by the Group of Experts on the Scientific Aspects of Marine Environmental Protection (GESAMP, 1989), much of which was subsequently published as a review article in the scientific literature (Duce et al., 1991). This paper briefly reviews results from several of these studies, especially those related to the Mediterranean Sea.

The biogeochemical effects of mineral dust deposition to the water surface are of interest from several perspectives. The observation of more than 30-fold increases in dust deposition during glacial periods suggests a link between climatic oscillations and carbon cycling in the surface ocean on global and glacial scales (Martin, 1990). This link is created through the roles played by eolian deposition in supplying Fe to remote parts of the open ocean and by iron limitation in controlling primary production. In these high-nutrient/low-chlorophyll (HNLC) regions, there may be a direct connection between atmospheric and biological processes (Martin, 1990; Kumar et al., 1995).

Dust input to the nutrient-limited Mediterranean basin ( $2.3 \times 10^6 \text{ km}^2$ ) is one of the greatest in the contemporary ocean ( $20\text{--}50 \times 10^6 \text{ tons yr}^{-1}$ ) and it is thus a natural laboratory in which to study the effects of dust deposition on the surface ocean in a setting that is distinct from the HNLC regions (Coale et al., 1996). There have been several hypotheses suggesting that this large dust input is responsible for biogeochemical effects, although no concerted scientific program aimed at elucidating these effects directly has yet been set up.

The aim of the present paper is to review current understanding of the magnitude and effects of dust inputs to the Mediterranean and to highlight important areas of uncertainty. Section 2 considers the magnitude of atmospheric dust inputs and problems related to their estimates. Section 3 compares fluxes of trace elements from atmospheric and riverine sources and considers evidence that atmospheric inputs affect the chemistry of surface waters in the Mediterranean. Section 4 looks at the role played by atmospheric processes in delivering nutrients to the surface Mediterranean and their spatial effects. Section 5 considers current hypotheses of mechanisms by which atmospheric deposition is believed to affect surface water biogeochemical cycles, focusing particularly on the roles of phosphate and iron, and reviews what evidence exists to evaluate them.

## **2. Magnitude and mineralogical composition of atmospheric mineral dust input**

Budgets and residence time estimates for trace metals and nutrients in the oceans have traditionally been based on riverine fluxes and have neglected atmospheric and other inputs. For most major and minor components of seawater, the atmosphere has generally been assumed to be a secondary source. However, over the last 20 years systematic studies of atmospheric inputs to the Mediterranean have been undertaken, and sufficient data are now available to allow fairly reliable estimates to be made. The most comprehensive study is that prepared by GESAMP (1989), together with several recent reviews devoted to the Western Mediterranean within the EROS 2000 project (Milliman & Martin, 1997).

For rivers, there is the major problem of understanding and quantifying transport through estuarine and coastal regions to deep waters. Milliman, Rutkowsky and Meybeck (1995) and GESAMP (1989) both attempted to address this problem by compiling run-off data for countries surrounding the Mediterranean. It is important to note, however, that many of these data reflect historic not present-day discharges. A good example of this is the decreased flow to the western Mediterranean, as illustrated in the introduction to the EROS-2000 DSR-II special issue (Milliman & Martin, 1997). The Rhone and many Italian rivers currently have discharges very similar to those of 40 years ago, whereas most Spanish and north African rivers have been dammed in the past 20–30 years.

Atmospheric flux estimates suffer from an undersampling problem. The atmosphere is a very dynamic compartment of the earth's system, within which concentrations of reactive gases and particulate matter vary considerably in space and time.

This variability ranges from rapid day-to-day changes in pollutant concentrations (lead, cadmium, NO<sub>x</sub>, SO<sub>x</sub>) to massive temporal and spatial changes in dust loading over the entire Mediterranean (Table 1), as a result of dust storms in North Africa (Guerzoni, Molinaroli & Chester, 1997; Bergametti et al., 1989; Dulac, Buat-Ménard, Ezat & Bergametti, 1989; Dulac et al., 1996; Loÿe-Pilot & Martin, 1996).

The high spatial and temporal variability of dust transport and deposition, coupled with the short duration of oceanographic research cruises, has made it difficult to obtain good estimates of dust deposition to those areas of the open ocean devoid of islands at which continuous sampling can be undertaken.

Encouragingly, however, recent results from ship-board collections of aerosol metals over the North Sea, where a steep spatial gradient in aerosol concentration is found (Yaaqub, Davies, Jickells & Miller, 1991), are comparable to samples collected at the coast (Chester, Murphy, Lin, Berry, Bradshaw & Corcoran, 1993). This finding suggests that atmospheric fluxes estimated from coastal sites may be extrapolated to adjacent marine areas. A similar result was found in the Western Mediterranean (WMED) where Martin, Elbaz-Poulichet, Guieu, Loÿe-Pilot and Han (1989) and Chester et al. (1993) showed that background aerosols at sea have similar Pb and Cd values to those collected at a remote coastal station in Sardinia (Guerzoni et al., 1997).

Despite the shortcomings of these estimates, for many trace elements (e.g., Al, Pb, Zn), it has been shown that the local dissolved flux from partial dissolution of eolian dust in surface waters is much greater than that carried by the Rhone and Po rivers into the WMED and Adriatic respectively (Guieu et al., 1997; Guerzoni, Molinaroli, Rosini, Rampazzo, Quarantotto & Cristini, 1999). Even where river fluxes appear to be similar in magnitude to those of dust, estuarine and near-shore removal processes attenuate river fluxes severely, often resulting in little of the riverine signal penetrating into offshore regions. An exception to this is Cd, which forms stable and soluble complexes with chlorides and may also be desorbed from river-borne particulate during estuarine mixing, so that in this case the input very often influences open basin waters (Boyle, Chapnick, Bai & Spivack, 1985; Elbaz-Poulichet et al., 1989).

Table 1

Range of insoluble particulate load concentrations in air and rain during two years at Sardinia (39°N; 9°E: from Guerzoni et al., 1997)

	Air ( $\mu\text{g m}^{-3}$ )	Rain ( $\text{mg l}^{-1}$ )
Background <sup>a</sup>	1–3	0.1–10
Desert-enriched <sup>b</sup>	20	10–50
Saharan outbreaks <sup>c</sup>	44	50–500
Maximum recorded	110	510

<sup>a</sup> Al/Si<0.3; Ca<2%; associated rain pH<5.60; Ca<200 mEq/l

<sup>b</sup> Al/Si>0.3; Ca>2%;

<sup>c</sup> Al>1000 ng m<sup>-3</sup>; Al/Si>0.4; Ca>4%.

2.1. Atmospheric mineral particle fluxes and mineralogy

Beside the undersampling problem, several theoretical problems also hinder our ability to derive atmospheric flux estimates to the ocean from atmospheric suspended loads. It is therefore useful to compare the measured data with the results of GESAMP flux calculations (Duce, 1991).

Fig. 1 and Table 2 list most of the coastal stations where long (2–11 yrs) time series data are available together with an inventory of mean annual bulk dust fluxes around the Mediterranean. Flux values are deduced from a combination of aerosol suspended loads, dry, wet and bulk deposition samples. Sampling frequency at the various sites ranged from a few days (aerosols) to a few weeks (bulk).

Mean annual deposition mass fluxes of mineral dust for the WMED range from 3–12 g m<sup>-2</sup> yr<sup>-1</sup>, whereas much higher values are found (20–50 g m<sup>-2</sup> yr<sup>-1</sup>) in the Eastern Mediterranean (EMED). The measured data are higher than those calculated with the GESAMP model using precipitation rates and a scavenging ratio (SR) of 200. As the SR for the Mediterranean is probably higher (Guerzoni, Quarantotto, Cesari, Molinaroli, Rampazzo & Le Bolloch, 1996), mass fluxes were recalculated using an SR of 500, and are listed in Table 2. When SR=500 is used, the two estimates agree very well.

With the data from Table 2 we can calculate mean values of 8, 12 and 35 g m<sup>-2</sup> yr<sup>-1</sup> for WMED, CMED and EMED respectively, and use these values to estimate the total annual atmospheric dust flux, which turned out to be ~40×10<sup>6</sup> tons for the whole Mediterranean basin (Table 3). This estimate may be compared to total riverine discharge estimates recently revised by LOICZ (Milliman et al., 1995).




Fig. 1. Arrows indicate the location of sampling sites for which time series of atmospheric data are available. Squares: sediment trap locations (B=Balear; D=Dyfamed; P=Paleoflux). Division in sub-areas 1-10 according to UNEP (1984) for budget calculations.

Table 2

Mean annual (and range) atmospheric mass fluxes ( $\text{g m}^{-2}$ ) measured at various coastal sites (refer to Fig. 1 for site locations). Mean fluxes for three sub-basins calculated by averaging Spain, SE France, Sardinia and Corsica for WMED, Corsica, NE Italy and Crete for CMED, and Crete and Israel for EMED. Estimates from GESAMP model (Duce et al., 1991) calculated with scavenging ratio values of 200 and 500 respectively

Site	Flux $\text{gm}^{-2}$	Sub-basin mean ( $\text{gm}^{-2}$ )		GESAMP SR=200 (500)
Spain <sup>a</sup>	5 (1–11)			
SE France <sup>b</sup>	4 (2–11)			
Sardinia <sup>c</sup>	8 (2–12)	WMED	8	3.4 (8)
Corsica <sup>d</sup>	12 (4–25)			
NE Italy <sup>e</sup>	3 (2–6)	CMED	12	7 (18)
Crete <sup>f</sup>	21 (6–46)			
Israel <sup>g</sup>	50 (36–72)	EMED	35	14 (35)

<sup>a</sup> Carratala, Bellot, Gomez and Millan (1996) and Roda, Bellot, Avila, Escarré, Pinol and Terradas (1993).

<sup>b</sup> Gueiu et al. (1997).

<sup>c</sup> Guerzoni et al. (1997); Le Bolloch and Guerzoni (1995).

<sup>d</sup> Loÿe-Pilot et al. (1986); Bergametti et al. (1989).

<sup>e</sup> Guerzoni et al. (1999); Rossini and Guerzoni (personal communication).

<sup>f</sup> Nihlen and Mattsson (1989) and Pye (1992).

<sup>g</sup> Herut and Krom (1996).

The sub-areas indicated by GESAMP (Nos. 1 to 10, shown in Fig. 1) were used for calculations.

### 2.1.1. Mineralogy

Loÿe-Pilot, Martin and Morelli (1986) and Guerzoni et al. (1997) have shown the relationship between eolian deposition and deep sea sedimentation, and found that the atmosphere contributes on average from 10 to 30% of recent sediments. Here, we highlight differences among sub-basins and the possible role of mineralogy in identifying the sites where the atmospheric input is most important.

The mineralogical signature may be used as an indicator of dust origin. Mineral dust mainly consists of a mixture of silicates (clay minerals, feldspar, quartz) associated with carbonates. The abundance of each of these minerals in dust is highly variable, mainly reflecting the source composition and its evolution during transport. Several studies have examined specific minerals within the clay to identify source regions. For example, palygorskite has been used as an indicator of dusts of north-west African origin in samples collected over the Mediterranean (Tomadin & Lenaz, 1989; Molinaroli & Ibbá, 1995; Molinaroli, 1996).

Another approach is to look for mineral associations which are characteristic of dust sources. For example, illite is an ubiquitous clay and is not characteristic of any particular weathering regime, whereas kaolinite, which may also be produced in a variety of weathering conditions, is characteristic of desert weathering regimes. A comparison of the mineralogy of dusts collected throughout the Mediterranean

Table 3

Comparison of riverine discharge (R) and atmospheric (A) input of insoluble particles to Mediterranean Sea. Data on riverine discharge are from Milliman et al. (1995). Atmospheric mass inputs derived from average fluxes in three sub-basins (west to east) of 8–12–35 g m<sup>-2</sup> yr<sup>-1</sup>, respectively (see Table 2 and text for explanation)

Sub-basin	Area (10 <sup>6</sup> km <sup>2</sup> )	Mass (10 <sup>6</sup> tons)		
		R	A	%A
WMED	0.84	30	7	19
CMED	0.80	35	10	22
EMED	0.70	10	24	70
Total	2.34	75 <sup>a</sup>	41	35

<sup>a</sup> if 2/3 of riverine input remains on the continental shelf (<100 m) then the offshore riverine discharge is reduced to 25×10<sup>6</sup> tons, and the atmospheric contribution increases to ~60%.

shows that illite reaches its highest value in the CMED and its lowest in the EMED (Table 4). In the CMED mixed dusts are observed. Kaolinite is more abundant in dust originating from the eastern compared to western Sahara, the highest concentrations being observed off the Egyptian coast. Smectite is very common in the EMED but not in the WMED. A comparison between dust and sediment has been attempted (Caqueneau, Gaudichet, Gomes, Magonthier & Chatenet, 1999) by considering the difference between illite and kaolinite distributions in the Mediterranean and the illite/kaolinite ratio in the African source dust. The comparison between the I/K ratio in dust and sediments is shown in Table 4. WMED dust transported from the Sahara originating from north-west Africa shows an I/K ratio of 1.8. Caqueneau et al. (1999) observed a range of I/K values from 2.4 to 1.6 in north-west Africa. The similarity of the I/K ratio between dust and sediment (I/K=1.8) in the WMED indicates that the major contribution to sedimentation in this region is from dust

Table 4

Average of clay mineral compositions and I/K ratio in Saharan dusts (D) and sediments (S) in MED basin<sup>a</sup>

Sample	Illite	Chlorite	Kaolinite	Smectite	I/K
WMED (D)	53	9	29	9	1.8
WMED (S)	48	10	26	16	1.8
CMED (D)	66	12	16	3	4.0
CMED (S)	45	12	13	30	3.5
EMED (D)	32	5	38	25	0.8
EMED (S)	15	5	35	45	0.4

<sup>a</sup> This average was compiled using data from several authors: Chester, Sharples, Sanders and Saydam (1984) (3 samples); Mazzucotelli et al. (1986) (3 samples); Molinaroli (1996) (10 samples) for the WMED. Lenaz et al. (1988) (5 samples) for the CMED. Chester, Baxter, Behairy, Connor, Cross, Elderfield et al. (1977) (7 samples); Ganor and Foner (1996) (17 samples); Tomadin and Lenaz (1989) (3 samples) for the south-EMED; Tomadin (1981) (3 samples) for the (S) WMED; Tomadin and Borghini (1987) (4 samples) for the (S) CMED; Venkatarathnam and Ryan (1971) (20 samples) for the (S) EMED.

deposition from the western Sahara. This conclusion contradicts data from Table 3, which indicate an overwhelming river contribution in this region. The fact that the sediment I/K ratio appears to be set by atmospheric dust probably indicates that a large fraction of the riverborne material is deposited on shelves.

The CMED has an average I/K ratio of 4.0 in dust and 3.5 in sediments. This very high ratio shows that the dust is a mixture of Saharan and background dusts (European), since Caquineau et al. (1999) demonstrated that the I/K ratio is always  $<2.4$  in African regions.

The EMED has a low I/K ratio (0.8) in dust and 0.4 in sediments. A relatively low I/K ratio (0.7) is associated with north east African sources by Caquineau et al. (1999). The similarity between dust and sediment values again indicates that the major contribution to sedimentation is from dust deposition, in particular from the eastern Sahara.

## 2.2. *Sediment trap estimates*

Many sediment trap studies have been or are being carried out in the Mediterranean, e.g. Mediterranean Targeted Projects 1 and 2 (MTP1 and MTP2), but very few of these are useful for studying dust deposition, mainly because of their location relative to river inputs.

For instance, among the traps in the WMED, very different results have been derived from those deployed in the Balearic basin ( $39^{\circ}\text{N}$ ,  $3^{\circ}\text{E}$ ; 'B' in Fig. 1) with mass fluxes of  $50\text{--}70\text{ g m}^{-2}\text{ yr}^{-1}$ , compared to the DYFAMED site ( $43^{\circ}\text{N}$ ,  $8^{\circ}\text{E}$ ; 'D' in Fig. 1) where average mass fluxes are  $31\text{--}35\text{ g m}^{-2}\text{ yr}^{-1}$ . In addition, the relative contribution of the lithogenic fraction is different between the two sites (60% and 25%, respectively). The high flux and lithogenic contribution to the Balearic site is probably related to bottom resuspension and coastal run-off.

The trap used in Paleoflux, MARFLUX ( $34^{\circ}\text{N}$ ,  $20^{\circ}\text{E}$ ; Ziveri, De Lange, Rossignol-Strick, Schrader & Thomson, 1996; 'P' in Fig. 1) is in the best location for dust flux measurement in the EMED, since river discharge is negligible in this region. The total annual average flux recorded in this 3000 m deep trap deployed between November 1991 and August 1994, was around  $25\text{ g m}^{-2}\text{ yr}^{-1}$ , with approximately 60% of lithogenic origin (Rutten, De Lange, Ziveri, Thomson, Van Santvoort, Colley et al., 1999; Ziveri et al., personal communication).

Heussner and Monaco (1996) showed that two flux gradients of total mass are evident in the Mediterranean, and that they are related to trophic conditions. A gradient of decreasing flux is seen on the scale of the entire Mediterranean basin from west to east, roughly corresponding to the degree of oligotrophy. Within each sub-basin, a second gradient of trophic conditions appears, of decreasing flux from north to south. The inorganic carbon flux is observed to be slightly higher in the EMED, perhaps related to desert-derived carbonate inputs. Heussner and Monaco (1996) conclude that seasonal variations in fluxes in the traps essentially result from changes in continental input rates to the Mediterranean system, i.e., direct deposition of dust through rain and the effect of rain-induced river run-off. The sensitivity of traps to river run-off explains why few of these sites can be used for absolute and relative

quantification of Saharan dust export flux to the Mediterranean. In fact, as far as we are aware, it has been difficult (if not impossible) to distinguish between lithogenic material from dust and that from river discharge in trap samples.

Despite these problems, trap data from the DYFAMED (WMED) and Paleoflux (EMED) sites appear to be promising. At both sites, the lithogenic flux appears to be similar to, or smaller than, the eolian fluxes estimated for the region from coastal sites (Table 2). It is very important that, in future trap work, wherever possible, sites that are suitable for the task of quantifying dust deposition are chosen. The EMED is a particularly important region that needs more coverage of eolian deposition from sediment traps.

### 2.3. *Remote sensing and modelling*

In a series of recent papers, Dulac et al. (1996) and Moulin, Guillard, Dulac and Lambert (1997a), presented an analysis of historical Meteosat data, which monitored dust export from Africa over the Mediterranean on a quasi-daily basis between June 1983 and December 1994. These authors showed that, despite large daily variability, there were clear seasonal cycles with minima during the dry season (November, December and January). The general pattern is that dust transport begins over the eastern basin in spring and spreads over the western basin in summer.

Tentative estimates of deposition fluxes were made from Meteosat vertically integrated dust concentrations, using simple deposition velocity approaches, on a yearly time scale at the DYFAMED station, and for shorter periods in the western basin (Dulac et al., 1996). A similar paper was published with results for the DYFAMED station in the Ligurian Sea (Moulin, Lambert, Dulac & Dayan, 1997b). The former authors compared Meteosat data with measured aerosols at the Corsica site for a six-month period (1/2/87–31/8/87) and found the average measured flux was  $0.024 \text{ g m}^{-2} \text{ d}^{-1}$  (equivalent to approximately  $9 \text{ g m}^{-2} \text{ yr}^{-1}$ ), whereas Meteosat data yielded a flux of  $0.01\text{--}0.03 \text{ g m}^{-2} \text{ d}^{-1}$  ( $4\text{--}11 \text{ g m}^{-2} \text{ yr}^{-1}$ ) for the same period.

Dulac et al. (1996) also compared Meteosat-derived atmospheric fluxes in 1987 and 1988 with seawater dust fluxes measured in sediment traps at 200 m water depth during the same period at the DYFAMED station. If the periods with sediment trap data are examined, there are results for 364 days over a 2-year period. Seasonal variations are shown in Fig. 2. The minimum and maximum Meteosat-derived fluxes bracket the trap fluxes in summer and autumn 1987 and again between spring and autumn 1988. In both winters (1987 and 1988) the trap flux is greater than the maximum Meteosat estimates.

The Meteosat estimates show a similar range for both years ( $3.5\text{--}10.8$  and  $3.6\text{--}11.0 \text{ g m}^{-2} \text{ yr}^{-1}$ ) while the sediment trap data show distinctly different fluxes ( $12$  and  $9 \text{ g m}^{-2} \text{ yr}^{-1}$ ) in the two years. Some of this variability may result from the various short-term hydrological and biological processes that control the downward transport of particulate material in the water column. The susceptibility of trap data to short-term variations is shown in Fig. 3, showing contemporaneous trap fluxes and Meteosat estimates.



ber' aerosols have strikingly different trace metal composition, and this is clear in samples collected on board vessels as well as at coastal stations (see Table 5).

Many studies have documented the airborne concentrations of trace metals over the Mediterranean Sea (Dulac, Buat-Ménard, Arnold & Ezat, 1987; Guerzoni, Lenaz & Quarantotto, 1988; Bergametti, Dutot, Buat-Ménard, Losno & Remoudaki, 1989; Chester, Nimmo, Murphy & Nicolas, 1990; Chester et al., 1993; Migon & Caccia, 1990; Mateu, Colom, Forteza & Cerda, 1993; Migon, Alleman, Leblond & Nicolas, 1993; Sandroni & Migon, 1997). Most metals have a relatively short residence time in the troposphere (between a few days and a few weeks: Bergametti, 1987; Buat-Ménard, 1993). They are not well mixed in the atmosphere and thus aerosols and rainwater are expected to exhibit strong spatial and temporal variability (Buat-Ménard, 1993). However, while seasonal variability is observed in the NW Mediterranean (Nicolas, Migon, Leblond & Journal, 1995), spatial variability is rather low for metals such as Al, Cd, Cu, Fe, Pb and Zn, and the annual averages from 7 coastal sampling sites are within the same order of magnitude (Table 6) (Chester et al., 1990, 1993; Guieu et al., 1997; Guerzoni et al., 1996; Sandroni & Migon, 1997).

### 3.2. Metal fluxes (total, dry, wet)

Several authors who have attempted to estimate the atmospheric fluxes of metals to the surface ocean (Chester, Nimmo & Murphy, 1989; Dulac et al., 1989; Guieu, Thomas, Martin & Brun-Cottan, 1991; Migon, Morelli, Nicolas & Copin-Montégut, 1991; Guerzoni, Landuzzi, Lenaz, Quarantotto, Rampazzo, Molinaroli et al., 1993; Nicolas et al., 1995; Migon, Journal & Nicolas, 1997) have pointed out the difficulties encountered in quantifying the different modes of atmospheric inputs, i.e. wet/dry/total and dissolved/particulate. While evaluation of wet inputs, calculated from rainwater concentration and rainfall amount, is generally reliable, that of dry deposition is more problematic. It has been calculated by a number of methods: i)

Table 5

Mean trace metal concentrations (conc. units:  $\text{ng m}^{-3}$  of air) and  $\text{EF}_{\text{crust}}$  values for European anthropogenic-dominated and Saharan crust-dominated particulate aerosol populations collected over Tyrrhenian Sea ( $n=9$ , Chester et al., 1984) and at Sardinia ( $n=55$ , Guerzoni et al., 1999)

	European anthropogenic-dominated aerosol population ( $n=5+48$ )		Saharan crust-dominated aerosol population ( $n=4+7$ )	
	Concentration ship-Sardinia	$\text{EF}_{\text{crust}}$ ship-Sardinia	Concentration ship-Sardinia	$\text{EF}_{\text{crust}}$ ship-Sardinia
Al	112–210	1.0–1.0	2000–2929	1.0–1.0
Fe	117–84	1.7–8.3	1249–1139	0.9–1.7
Mn	4.3–5.4	3.5–1.5	19–24	0.8–1.1
Cr	1.0–0.5	8.6–3.3	4.9–3.3	2.9–1.1
Ni	1.3–5.3	15–21	3.1–7.3	2.8–4.1
Zn	12–22	120–63	8–28	6–18
Pb	11–12	767–550	7–25	46–66
Cd	0.44–0.17	1219–80	0.31–0.08	96–12

Table 6  
Average trace metal concentrations and  $EF_{crust}$  values for particulate aerosols from a number of Mediterranean Sea sites (conc. units:  $ng\ m^{-3}$  of air)

	Spain <sup>a</sup> Conc.	$EF_{crust}$	Corsica <sup>b</sup> Conc.	$EF_{crust}$	Sardinia <sup>c</sup> Conc.	$EF_{crust}$	Erdemli <sup>d</sup>	Cap Ferrat <sup>e</sup> Conc.	$EF_{crust}$	Vignola <sup>f</sup> Conc.	$EF_{crust}$
Al	398	1	168	1	480	1	680	370	1	109	1
Fe	316	1.3	144	1.25	278	0.85	685	320	1.3	–	–
Mn	10	2.2	4.3	2.7	7.4	1.3	12.6	11	262	1.7	1.4
Cr	1.8	3.4	–	–	0.51	0.89	10.8	2.5	5.6	–	–
Cu	7.9	29	2.1	18	–	–	–	6.2	24	1.9	26
Zn	50	148	19	133	21	52	19	41	130	12	130
Pb	50	837	16	635	14	194	30	58	1045	9	550
Cd	0.60	628	0.66	1633	0.30	260	0.19	0.60	676	0.11	423

<sup>a</sup> Chester et al. (1993).

<sup>b</sup> Bergametti et al. (1989).

<sup>c</sup> Keyse (1995), Guerzoni et al. (1996).

<sup>d</sup> Kubilay and Saydam (1995).

<sup>e</sup> Chester et al. (1990).

<sup>f</sup> Migon et al. (1993).

on the basis of experimental cascade impactor data (e.g., Bergametti, 1987; Remoudaki, 1990); ii) from theoretical models (e.g., Dulac et al., 1989); and iii) with surrogate surfaces (e.g., Dolske & Gatz, 1985; Baeyens, Dehairs & Dedeurwaerder, 1990; Guieu, 1991). Results differ greatly according to methodology. Comparisons between estimates must be interpreted with caution, principally as a result of poor knowledge of elemental mass-size distributions (Dulac et al., 1989; Migon et al., 1991; Buat-Ménard, 1993). For example, in the NW Mediterranean, Dulac et al. (1989) proposed a theoretical dry deposition rate of  $0.04 \text{ cm s}^{-1}$  for Pb, while Remoudaki (1990) gave an experimental value of  $1.9 \text{ cm s}^{-1}$ . Dry deposition rates are poorly known for submicrometer particles (Slinn, 1983; Arimoto & Duce, 1986), and difficulties increase when particle size exceeds  $10 \mu\text{m}$ : in this case, deposition rates are very high and their estimation becomes very questionable (Duce et al., 1982). Moreover, phenomena such as fog or dew, which are common in coastal areas, significantly enlarge particles, thereby increasing their deposition rate (Bergametti, 1987). Despite these problems, there is a significant amount of data on trace metal atmospheric fluxes, mainly for the Western and Central Mediterranean. They are listed in Table 7, which shows that the range of fluxes is quite high for some metals (e.g., Al, Cd, Zn), and lower for others (Cu, Pb, Fe, Ni) (Guieu et al., 1997; Guerzoni et al., 1999).

### 3.3. Fate of trace metals in seawater

The magnitude of air-to-sea fluxes of trace metals partly depends on their concentrations in the air. However, the fate of an atmospherically-transported trace metal, once deposited at the sea surface, depends on its thermodynamic speciation in seawater, its speciation in the parent aerosol, and the kinetics controlling any speciation change. Aerosol speciation may be assessed by determining the partitioning of trace metals between the following fractions: (i) exchangeable, the most mobile, (ii) oxide/carbonate, intermediate mobility, and (iii) refractory, the least mobile. The speciation of trace metals in aerosols is a function of the source of the aerosol since some trace metals have very different speciation signatures in anthropogenic-rich

Table 7  
Range of total atmospheric fluxes (insoluble and soluble fractions), calculated from wet and dry deposition

$\mu\text{moles m}^{-2} \text{ yr}^{-1}$	CMED <sup>a</sup>		NWMED <sup>b</sup>	
	insoluble	soluble	insoluble	soluble
Al	11250–33000	330–890	4900–31500	370–3780
Cd	0.1–0.3	0.5–2.8	0.9–3.6	1.8–8.0
Cu	50–70	45–72	22–30	12–20
Fe	2600–7300	90–180	12500–14500	570–2500
Ni	8–32	17–24	7–9	7–9
Pb	5–38	5–24	6–11	4–9
Zn	46–76	210–460	15–1000	15–1150

<sup>a</sup> Guerzoni et al. (1999).

<sup>b</sup> Guieu et al. (1997).

and crust-rich ‘end-member’ Mediterranean aerosols (Chester, Nimmo & Keyse, 1996), and chemical changes occur during passage and transformation in the atmosphere prior to deposition on the sea surface.

Al and Fe are generally refractory in both end-member aerosols, Mn is speciated between all three fractions in both end-member aerosols, but Cu, Zn and Pb switch speciation signatures between the two end-member aerosols, being essentially refractory in crustal aerosols and exchangeable (i.e., mobile) in anthropogenic ones. Aerosols are removed from the air by dry fall-out and wet precipitation scavenging deposition modes.

In the dry mode, aerosols reach the sea surface directly and trace metal solubility is constrained by aerosol-sea water reactivity. There is a direct relationship between the extent to which a trace metal is soluble in seawater and the extent to which it is held in the exchangeable fraction of an aerosol (Fig. 4). For Al and Fe in both end-member aerosols,  $<10\%$  of their total concentrations is in the form of exchangeable associations and both are also relatively insoluble in seawater. However, since both metals are present in relatively high concentrations in aerosols (3–8% by weight), even a solubility of a few percent can release considerable quantities of Al and Fe to seawater in a dissolved form. Mn has between  $\sim 20\%$  and  $\sim 50\%$  of its total concentration in the exchangeable fraction of each end-member aerosol and is relatively soluble, with  $\sim 35\%$ – $\sim 50\%$  of the total amount from each phase dissolving. Cu, Zn and Pb have significantly higher solubilities from the anthropogenic-rich than from the crust-rich end-member, because of their different speciation signatures in the two aerosol types. This is important, because it means that the aerosol seawater solubilities of Cu, Zn and Pb are reduced when crust-rich Saharan inputs perturb the anthropogenic-rich background aerosol over the Mediterranean.

In the wet deposition mode, trace metal solubility is constrained initially by aerosol-rainwater reactivity since some metals are already in solution prior to the deposition of the aerosols on the sea surface. Solubility is still a function of trace metal speciation in the parent aerosol, as illustrated by the dissolved/particulate speciation




Fig. 4. Relationship between percentages of total concentrations of trace metals soluble in seawater and in exchangeable associations in A) anthropogenic-dominated and B) crust-dominated aerosols. After Chester et al. (1993).

of a number of trace metals in rainwater from Western Mediterranean sites (Table 8). Al and Fe are generally insoluble in all types of rainwater (<20% of total concentrations), and Cd and Mn are relatively soluble (60–90% of total aerosol). Cu, Zn and Pb solubilities vary considerably from one rainwater sample to another and from site to site, e.g. that of Pb varies from ~20% at Sardinia to ~70% at Cap Ferrat.

The speciation and solubility of trace metals is strongly affected by rainwater pH. Scavenging of anthropogenic-rich aerosols results in rainwater having pH values of <~4–5. In contrast, the scavenging of Saharan dust, with relatively high carbonate content, yields rainwaters with pH values of >~5. Many adsorption/desorption reactions are pH-dependent, particularly those affecting exchangeable trace metals. This may be illustrated for Pb in a series of rainwaters collected at Cap Ferrat (Chester et al., 1990). The most striking feature in the pH vs solubility plot for Pb is the presence of the classical pH adsorption edge (Fig. 5), which is typical of adsorption/desorption processes at particulate-water interfaces and has been studied by various authors, in both rain and aerosols (Chester et al., 1993; Lim, Jickells, Colin & Losno, 1994; Spokes & Jickells, 1996). For Pb, this adsorption edge lies in the pH region ~4.8–~5.8. On the more acidic side, ~80% of total Pb is in the soluble form, whereas on the alkaline side only ~10% is soluble. However, some authors note that the low solubility region is defined by a relatively small number of points, reflecting the few cases studied of rain with pH >6 (Lim et al., 1994; Spokes & Jickells, 1996). This observation is important, given that these waters normally contain variable amounts of particulate material and that particle concentrations, grain-size, and physical surface characteristics (scavenged clays or silicate mineral aerosol) may also be important parameters in adsorption models of trace metals in rain (Lim et al., 1994). In fact, aerosol dissolution experiments performed at particle concentrations ranging from 3 to 300 mg l<sup>-1</sup> (the usual particle concentrations found in Mediterranean rainwater, see Table 1) showed that Cd and Pb have

Table 8

Average dissolved/particulate speciation of trace metals in rainwaters from a number of Mediterranean Sea sites; data given as % of total trace metal concentration in dissolved phase

	Cap Ferrat <sup>a</sup>	Tour du Valat <sup>a</sup>	Sardinia <sup>b</sup>	Corsica <sup>a</sup>
Al	18	19	2	8
Fe	–	11	3	13
Mn	60	63	–	67
Ni	54	58	–	–
Co	61	50	–	–
Cu	82	71	–	49
Zn	–	68	–	76
Pb	65	52	21	48
Cd	92	75	75	–

<sup>a</sup> Guieu et al. (1997).

<sup>b</sup> Guerzoni et al. (1997).




Fig. 5. pH solubility of Pb in Western Mediterranean rainwaters (after Chester et al., 1990).

low solubility at high particle concentrations (Fig. 6). The solubility of Pb was also found to increase with increasing  $EF_{\text{crust}}$  values and with increasing content of the finest ( $<1 \mu\text{m}$ ) grain-size of the aerosols (Guerzoni et al., 1999).

### 3.4. Atmospheric versus riverine inputs

Having identified atmospheric input as being important to the ocean, what problems remain in understanding the magnitude and significance of this input mode, with respect to riverine input? Aerosol undersampling problems may be remedied by increased sampling effort. Alternatively, atmospheric fluxes may be calculated from atmospheric transport and deposition models (Erdman, Sofiev, Subbotin, Dedkova, Afinogenova, Chesukina et al., 1994), using information on emission source strength (Nriagu & Pacyna, 1988; GESAMP, 1989; Axenfeld, Munch, Pacyna, Duiser & Veldt, 1992).

We present here two examples of comparison between riverine and atmospheric inputs using measurements made in the WMED and the Adriatic, Central Mediterranean (CMED).

#### 3.4.1. Measurements

The total atmospheric inputs of several trace elements and potential pollutants are compared with total riverine trace metal inputs for the Adriatic and the NW Mediterranean in Table 9. The data indicate that the atmosphere supplies 16–90% of the total input of these elements. When dissolved fluxes are compared (the form most likely to impinge on surface water biogeochemical cycles), the influence of atmospheric input is even more pronounced (Table 10). For example the atmosphere is 5–50 times more important than rivers for dissolved Zn fluxes and 15–30 times more important for Pb fluxes. A similar pattern is seen for Cu. The dramatic difference between total particulate and dissolved fluxes underscores the important role played




Fig. 6. Lead solubility versus particulate load in aerosol dissolution laboratory experiment (from Guerzoni et al., 1999).

by speciation of trace elements in rainwater in determining their flux to the surface ocean.

### 3.4.2. Models

Only by using models can we extend the same comparison to each of the three basins of the Mediterranean (Table 11). This comparison is meant to show the same approach as the two tables above but is extended to other basins for total fluxes.

Available estimates of riverine and atmospheric inputs were computed on the basis of average pollutant concentrations and mean water discharges for rivers (UNEP,

Table 9

Comparison between total inputs of atmospheric and riverine origin for Adriatic and NW Mediterranean (all data in ton yr<sup>-1</sup>)

	tons year <sup>-1</sup>	Al <sup>a</sup>	Fe <sup>a</sup>	Cd	Pb	Cu	Ni	Zn
Adriatic <sup>b</sup>	atmosphere	29	14	18	311	489	148	177
	Po river	86	74	7	151	281	514	935
NW Med. <sup>c</sup>	atmosphere	97	151	126	579	486	162	12,600
	Rhone and Ebro rivers	441	232	13	318	505	450	1545

<sup>a</sup> Al and Fe: ton×1000

<sup>b</sup> Adriatic Sea area: 112,000 km<sup>2</sup>; from Rossini and Guerzoni (personal communication)

<sup>c</sup> NW Mediterranean Sea area: 180,000 Km<sup>2</sup>, from Guieu et al. (1997)

Table 10

Comparison between dissolved inputs of atmospheric and riverine origin for Adriatic (area: 112,000 km<sup>2</sup>) and NW Mediterranean (area: 180,000 Km<sup>2</sup>)

		tons year <sup>-1</sup>	Al	Fe	Cd	Pb	Cu	Ni	Zn	PAH	PCB
Adriatic <sup>a</sup>	atmosphere	1039	573	13	152	332	116	1560			
	Po river	60	30	4	10	66	148	215			
NW Med. <sup>b</sup>	atmosphere	10	16	99	252	189	81	6813	45	12	
	Rhône and Ebro rivers	1	1	2	8	175	115	105	15	1	

<sup>a</sup> Adriatic Sea: from Rossini and Guerzoni (personal communication).

<sup>b</sup> NW Mediterranean Sea: from Guieu et al. (1997).

1984) and by applying a Eulerian–Lagrangian model for atmospheric transport and deposition (Erdman et al., 1994).

Several sources of errors are associated with these estimates. An example is given in Table 12, showing a comparison between measured and model data for Pb and Zn in the Adriatic (equivalent to WMO area MT5; see Erdman et al., 1994) and north-western Mediterranean (equivalent to WMO area MT2; see Erdman et al., 1994). In these models, the authors state that the discrepancy between regional mean values for Pb does not exceed 20%, but very often individual data are less reliable. For Zn, the total underestimation of mean values by the models may be 10 times lower, mainly because of underestimation of the emission strength.

These comparisons of modelling results with measurement data for atmospheric deposition should only be considered as approximate, for the following reasons (EMEP, 1994): i) models give grid square (e.g. 150×150 km) average values and measurements give point values, and variations between measurement and model data increase with increasing grid size and decreasing averaging time; ii) as completeness of measurement data should be 75–80% over a rather long period of time (e.g., at least one month) to be used for comparison with modelling results, better spatial and temporal data coverage is needed; iii) measurement sites should be representative for the whole grid square (meteorological conditions, surface character-

Table 11

Comparison of total riverine (R) and atmospheric (A) inputs to Mediterranean Sea (all in tons yr<sup>-1</sup>). Data on Pb and Zn riverine inputs from UNEP (1984), on atmospheric input from Erdman et al. (1994) and GESAMP (1989)

Sub-basin	Area (10 <sup>6</sup> km <sup>2</sup> )	Pb		Zn	
		R	A	R	A
Western Med	0.84	2200	3072	9200	1125
Central Med	0.80	1770	2757	10,700	809
Eastern Med	0.70	850	1605	4800	590
Total	2.34	4820	7404	24,700	2524
% atmosphere			61		9

Table 12

Comparison between atmospheric (measured and modelled) and riverine inputs (measured and estimated)

Region	Input		Pb tons y <sup>-1</sup>	Zn tons y <sup>-1</sup>	Reference
Adriatic	atmospheric	measured	311	177	Rossini and Guerzoni (pers. com)
		model	871	246	Erdman et al. (1994)
	riverine	Po river			
		measured	151	935	Rossini and Guerzoni (pers. com)
NW Med	atmospheric	estimated	1300	7900	UNEP (1984)
		measured	579	12600	Guieu et al. (1997)
		model	999	337	Erdman et al. (1994)
	riverine	Rhône and Ebro			
		measured	318	1545	Guieu et al. (1997)
	estimated	800	2400	UNEP (1984)	

istics, local pollution sources, etc.); iv) several monitoring sites are needed for a square of 150×150 km and correlation analysis between sites should be applied, taking into account seasonality, mixing, time scale, meteorology, emission and topography; v) measurement and modelling data used for comparison purposes are sometimes made at different times (or even in different years) and the year-to-year variability of meteorological conditions introduces uncertainty into single-year model (or measurement) estimates; vi) emissions used for modelling are rather uncertain and sometimes underestimated, and are mainly based only on annual estimates from major sources without the required temporal and spatial distributions.

It may be seen that atmospheric data are in agreement within a factor of 2–3, whereas huge differences are evident for riverine inputs. Some of these differences are the result of the reductions in run-off and discharge that have occurred over the last 10–20 years, and not always accounted for in the models. Furthermore, riverine model data are derived from run-off estimates and contain few real measurements, therefore providing only an order-of-magnitude estimate of the elemental input for the Mediterranean basin. In addition, the differences between models and measured data may be because of the behaviour of elements in estuarine areas. The chemical behaviour of elements in this zone is very difficult to generalise. For some elements there is removal from the water column in the estuarine zone while for others there is no effect. In some cases (e.g. Cd), there is addition of the element to the dissolved phase as a result of the competitive desorption of Cd bound to the surface of suspended particulate material by seawater ions. While the behaviour of a particular element can be characterised in a particular estuary, different physico-chemical regimes (e.g. pH, ionic strength, presence of dissolved organic matter, etc.) in a different estuary can lead to different behaviour in the mixing zone.

### 3.5. Case studies

#### 3.5.1. The atmospheric 'signature' in seawater

For some reactive trace elements including Al, Mn and most notably Pb (Bruland, Donat & Hutchins, 1991; Buat-Ménard, 1993; Patterson & Settle, 1987), the shapes of dissolved oceanic profiles with surface water concentration maxima suggest that atmospheric input is of major importance. For other elements which are not so rapidly removed from seawater by scavenging processes, internal biogeochemical cycles mask such input signals.

Al and Fe. It has been shown above that Al and Fe have only relatively low solubilities from aerosols but, since they are present in large concentrations, even low solubility can deliver significant quantities to the dissolved seawater pool from dust pulses.

The effect of dust deposition of Al on Mediterranean surface waters is dramatically visible in its effect on the surface water concentrations. The extremely high concentration of this element in crustal materials (8.1%; Taylor, 1964), and its low concentration in seawater makes it a particularly good tracer of the partial dissolution of eolian dust in Mediterranean surface waters. In fact, as a result of biological and convective transport, the deep waters of the Mediterranean have the highest concentrations of dissolved Al of any basin in the world, consistent with their receiving the greatest rate of deposition of eolian material. The effect that this has on the Al budget has been recognised by several workers (Hydes, De Lange & Debaar, 1988; Measures & Edmond, 1988; Chou & Wollast, 1997), the latter indicating that the partial deposition of eolian dust in the WMED was responsible for some 20 times as much input of Al as that supplied by rivers. That the eolian input is basin-wide is shown by the progressive build-up of Al from west to east along the advective flow path of the incoming Atlantic surface water (Fig. 7). In this figure, all post-1980 surface water Al values in the open Mediterranean are plotted as a function of longitude. Data from the region of the Gulf of Lions are excluded, to eliminate the local influence of the Rhone discharge. In addition, early Corsican nearshore data are excluded, to eliminate local artifacts.

The data distribution is quite distinctive in that there is an unmistakable increase in Al concentrations from 9–13 nM at the Strait of Gibraltar (6°W) to ~75 nM at 26°E in the eastern basin. The anomalously high value at ~5°E (100 nM) results from increased deep water mixing into the surface waters during December 1988, during a convection episode in the Western Mediterranean Deep Water formation region. The general trend indicates something of the order of a 60-nM Al enrichment of inflowing Atlantic surface waters during their eastward transit from the Strait of Gibraltar to the eastern basin.

The magnitude of this Al enrichment may be compared with estimates of atmospheric deposition to the region. Guieu et al. (1997), using atmospheric data from coastal collection sites, estimated an annual deposition of some 0.38 to 3.8 mmoles of dissolved Al m<sup>-2</sup> to the surface waters of the north-west Mediterranean. Such an input into a 50-m mixed layer, over the advective flow path of the Atlantic water, could easily raise Al concentrations in inflowing surface water to the values seen in




Fig. 7. Distributions of dissolved Al and Fe in surface waters of Mediterranean.

the eastern basin over the time-scales of surface water renewal in the Mediterranean (1–10 years). While the impact of dust is readily observed and is quantified in the Al signal, its effect on Fe concentrations is more difficult to quantify. The largest problem is the small data base of reliable measurements for Fe in the Mediterranean. Fig. 7 plots all the surface water dissolved Fe data of Morley, Burton, Tankere and Martin (1997) available in the EROS-2000 data base which appear to be oceanographically consistent. The data of Morley et al. (1997) show a wide range of concentrations from 0.11 nM in the Strait of Gibraltar to 2.83 nM south of the Gulf of Lions. However, unlike the Al, there is no clear geographic trend associated with these values. Nevertheless, it may be stated that, on the basis of this data set, Fe levels are generally above 1 nM within the Mediterranean. It therefore seems reasonable to conclude that Fe cannot be considered as limiting factor to phytoplankton. This appears to be probable merely on the basis of available nutrients. Since the surface Mediterranean is low in nutrients, Fe demand is similarly low. Using Fe limiting conditions and assuming a C:N ratio of 6.7, 0.5 nM of Fe (at the low end of the observed amount of Fe in 1-L of seawater) is capable of removing 7.5  $\mu\text{M}$  of N from surface waters, which is an extremely large amount of N relative to the concentrations of N normally occurring in 1-L of seawater in this nutrient-impooverished sea.

In summary, dust input to the Mediterranean clearly affects surface water trace element budgets, and this is most clearly seen for Al for which a large data base of modern observations exists. Although surface water Fe concentrations in the Mediterranean also appear high, the small size and geographically restricted existing data base precludes direct association of these values with dust deposition events. It is unlikely that dust deposition to the surface Mediterranean would spur primary pro-

duction through either the alleviation of Fe-limiting conditions or the provision of a bioactively reduced Fe species.

### 3.5.2. *Pb in atmosphere and seawater*

The EROS 2000 data base on lead concentrations over the period 1989–1993 gave an average of 100pM in surface waters and 80 pM in bottom waters. A mass balance for the WMED has been computed (Milliman & Martin, 1997), and the output/input ratio for lead in the whole basin is 0.8, with a residence time of 5.2 years (water residence time being ~15 years). A tentative budget of dissolved lead in the WMED (Wollast, pers. comm.) indicates an accumulation in the water of 440 tons yr<sup>-1</sup> for the period 1970–1985, related to the constant increment in consumption of leaded gasoline in that period.

In contrast, a recent paper (Migon & Nicolas, 1998) shows a reduction (over the period 1987–1995) of atmospheric fluxes of dissolved Pb from 3.5 to 1.5 mg m<sup>-2</sup> yr<sup>-1</sup>, and of surface water concentrations in the WMED (central Ligurian Sea) from 300 to 120 pM. The amount of Pb in inflowing rivers has probably decreased in a similar way to the atmospheric input. However: i) the natural Pb component is probably higher in rivers than in the atmosphere, because of soil erosion; ii) a previous study (Migon, 1993) showed that riverine Pb fluxes are far lower than atmospheric ones, and the loss of atmospheric Pb was estimated at >90%; iii) Pb dissolved in fresh water tends to precipitate and settle close to the river mouth (Sharp & Nardi, 1987; Windom, Smith, Rawlinsong, Hungspreugs, Dharmvanij & Wattayakorn, 1988; Dorten, Elbaz-Poulichet, Mart & Martin, 1991), which lowers the role of the riverine contribution to offshore marine Pb concentrations. These results confirm the importance of the atmosphere in delivering Pb to surface waters and in controlling its concentration. Fig. 8 illustrates the temporal evolution of dissolved Pb concentrations in surface waters (upper 100 m), together with mean atmospheric fluxes. The two curves have the same shape, which a posteriori indicates that the introduction of automotive Pb into the marine environment essentially occurs through the atmospheric pathway.

## 4. Atmospheric input of nutrients: estimates and impacts

The importance of the atmosphere as a source of nutrients, especially nitrogen, for terrestrial ecosystems was recognised as a by-product of studies of the impact of acid rain in the early 1980s. At the end of that decade, atmospheric nitrogen deposition to marine systems began to be investigated and its importance for the global ocean was assessed (Duce, 1986). For example, it has been noted that the atmospheric input of nitrogen may be partly responsible for the eutrophication of estuarine or coastal waters and the appearance of harmful algal blooms, by either direct input or through the watershed (Fisher, Ceraso, Mathew & Oppenheimer, 1988; Mallin, Paerl, Rudek & Bates, 1993; Paerl, Fogel & Bates, 1993).

Despite early observations of P limitation (Fiala, Cahet, Jacques, Neveux & Panouse, 1976; Berland, Bonin, Maestrini & Pointier, 1973; Berland, Bonin & Maestrini,




Fig. 8. Pb atmospheric flux and mean marine Pb concentrations (0-100 m) (from Migon & Nicolas, 1998).

1980), recent works suggest that Mediterranean surface waters are nitrogen-limited (Andersen & Nival, 1988; Owens, Rees, Woodward & Mantoura, 1989). In fact, there is growing evidence that the EMED is phosphorus-limited (Krom, Kress, Brenner & Gordon, 1991) and that the WMED is probably N-limited (Raimbault & Coste, 1990; Thingstad & Rassulzadegan, 1995), or that limitation shifts from nitrogen to phosphorus and vice versa depending on the period of the year (Fiala et al., 1976; Dolan, Thingstad & Rassoulzadegan, 1995) or the area considered (Woodward & Owens, 1989). The uniquely high levels of N/P ratios in the Mediterranean (20/27) compared with other open ocean averages (15) may reflect this situation, and are probably evidence of P limitation.

#### 4.1. Atmospheric inputs of nitrogen to the Mediterranean

While there are many natural sources of fixed nitrogen to the atmosphere —  $\text{NO}_3$  from soils and  $\text{NH}_3$  from wetlands — in the northern hemisphere the main sources are anthropogenic ( $\text{NO}_x$  from industrial combustion and vehicle traffic,  $\text{HNO}_3$  from vehicle traffic,  $\text{NH}_3$  emissions from intensive farming). The most abundant data are for inorganic wet deposition. Data for the WMED and EMED are listed in Table 13.

##### 4.1.1. Atmospheric input of wet inorganic nitrogen

The average wet dissolved inorganic nitrogen (DIN) flux to the north-west Mediterranean of  $\sim 100 \mu\text{mol N m}^{-2} \text{d}^{-1}$  (Lojze-Pilot, Martin & Morelli, 1990a) is somewhat higher than that to the south-east Mediterranean, which is  $\sim 60 \mu\text{mol N m}^{-2}$

Table 13  
Dissolved inorganic nitrogen (DIN) wet deposition

Studied site	Field data		Best estimates in sub-basin			Mediterranean	
	Period	Range (annual basis) $\mu\text{mol}$ $\text{m}^{-2} \text{d}^{-1}$	NWMED	WMED	EMED	$\mu\text{mol}$ $\text{m}^{-2} \text{d}^{-1}$	$10^{10}$ $\mu\text{mol}$ $\text{yr}^{-1}$
French coast — Camargue <sup>a</sup>	Feb 88/Jul 89	73					
French coast — Cap Ferrat <sup>b</sup>	86/87	119/142					2,5/7,5
Corsica — Bavella <sup>c</sup>	84/86	126/150	100	80	50	50/80	4,6/7,3
Sardinia — C.Carbonara <sup>d</sup>	91/94	47					
Sardinia — Torregrande <sup>e</sup>	92/93	41/56					
Israel — Haifa+South-EMED <sup>f</sup>	92/95	65			59	72	6,1

<sup>a</sup> Loÿe-Pilot et al. (1990a).

<sup>b</sup> Migon et al. (1989).

<sup>c</sup> Loÿe-Pilot et al. (1990b).

<sup>d</sup> Guerzoni et al. (1993).

<sup>e</sup> Le Bolloch and Guerzoni (1995).

<sup>f</sup> Herut and Krom (1996).

$\text{d}^{-1}$  (Herut & Krom, 1996), the value for the whole Mediterranean being between 50 and 80  $\mu\text{mol N m}^{-2} \text{d}^{-1}$  (Loÿe-Pilot et al., 1990a). Nitrate and ammonium contribute almost equally to the DIN, while nitrite is normally less than 3%.

#### 4.1.2. Atmospheric input of dry inorganic nitrogen

Dry deposition of inorganic nitrogen includes gas and particulate deposition. Only annual average data are available for atmospheric particulate inorganic nitrogen species. Values in the NWMED are 1.71 and 0.85  $\mu\text{g m}^{-3}$  for  $\text{NO}_3^-$  and  $\text{NH}_4^+$ , respectively (Klein, 1998), and in the EMED 1.51 and 0.97  $\mu\text{g m}^{-3}$  for  $\text{NO}_3^-$  and  $\text{NH}_4^+$  (Mihalopoulos, Stephanou, Kanakidou, Pilitsidis & Bousquet, 1997). These coastal values are in good agreement with the mean concentrations of 1.45 and 1.22  $\mu\text{g m}^{-3}$  for  $\text{NO}_3^-$  and  $\text{NH}_4^+$  obtained by Medinets (1996) from shipboard sampling over the whole Mediterranean. The reported values for particulate dry deposition depend on the deposition velocities chosen for  $\text{NO}_3^-$  and  $\text{NH}_4^+$ , the best estimate for the NWMED being 25  $\mu\text{mol N m}^{-2} \text{d}^{-1}$  (Klein, 1998).

Gas concentration measurements are very scarce, especially for  $\text{NH}_3$  which is assumed to be negligible regarding N deposition to the Mediterranean. The few  $\text{HNO}_{3\text{gas}}$  measurements are from urban sites.

Fig. 9 shows the partitions of various inorganic species and forms in the atmospheric deposition of nitrogen for the NWMED.

#### 4.1.3. Total inorganic nitrogen deposition (Table 14)

If only particulate dry species are taken into account, nitrogen dry deposition represents 25–33% of wet deposition in the north-west basin (Loÿe-Pilot, Martin & Mor-

elli, 1990b; Klein, 1998); taking  $\text{HNO}_{3\text{gas}}$  into consideration would increase the influence of dry deposition to equal wet deposition in coastal urbanised zones. Dry deposition equal to half that of wet deposition seems to be a realistic assessment for the open sea. Results are summarised in Table 14, which also lists data from the UNEP Mediterranean Pollution (MED POL) Project (computed from models) for 1991 (Erdman et al., 1994; UNEP, 1984). Total inorganic nitrogen deposition varies between  $214 \mu\text{mol N m}^{-2} \text{d}^{-1}$  for the Adriatic to about  $48 \mu\text{mol N m}^{-2} \text{d}^{-1}$  for the Central Mediterranean (UNEP zone VII) (Erdman et al., 1994), agreeing well with the previously estimated values of  $105 \mu\text{mol N m}^{-2} \text{d}^{-1}$  for the WMED and  $65 \mu\text{mol N m}^{-2} \text{d}^{-1}$  for the EMED (Loÿe-Pilot et al., 1990b).

Distance from the source and climatological conditions appear to be the major factors which control nitrogen inputs. If the DIN flux to the southern oceans,  $\sim 6 \mu\text{mol N m}^{-2} \text{d}^{-1}$ , represents the natural background level, then 90% of the total DIN flux into the Mediterranean is of anthropogenic origin.

4.1.4. Organic nitrogen

The importance of organic nitrogen in total atmospheric nitrogen deposition has recently been re-evaluated; it may represent half of the input of inorganic forms on a regional scale and be equal to them for global ocean (Cornell, Rendell & Jickells, 1995; Duce, 1997). The source of organic nitrogen is not known, but Cornell et al. (1995) believe that a large fraction of it is anthropogenic.

The few measurements of organic nitrogen content in the Mediterranean atmos-




Fig. 9. Relative importance of atmospheric inputs of different N species to NW Mediterranean; w=wet dp=dry particulate, dg=dry gaseous.

Table 14  
Total nitrogen deposition

	Best estimates				
	NWMED $\mu\text{mol m}^{-2} \text{d}^{-1}$	WMED $\mu\text{mol m}^{-2} \text{d}^{-1}$	EMED $\mu\text{mol m}^{-2} \text{d}^{-1}$	Mediterranean $\mu\text{mol m}^{-2} \text{d}^{-1}$	$10^{10} \text{ mol yr}^{-1}$
Field data <sup>c</sup>					
dry <sub>p</sub> =1/3 wet <sup>a,b</sup>	125	105	65	65/105	5,6/9,0
dry <sub>p+g</sub> =1/2 wet <sup>c</sup>	150	120	75	75/120	6,4/10,2
Model					
(UNEP MAP 94) <sup>d</sup>	117	107	80 without Adriatic 69	89	7,6

<sup>a</sup> Loÿe-Pilot et al., 1993.

<sup>b</sup> Klein, 1998.

<sup>c</sup> Loÿe-Pilot, 1998.

<sup>d</sup> Erdman et al., 1994).

<sup>e</sup> p=particulate, g=gaseous species.

phere (Spitzky, Ludwig & Wobig, 1990; Loÿe-Pilot, Cauwet, Spitzky & Martin, 1992) indicate that total amino acid concentrations in aerosols and rainwater are small (<5%) relative to inorganic species. It is clear that further studies are needed to assess the content, speciation and bioavailability of organic nitrogen in general and in the Mediterranean atmosphere in particular.

#### 4.1.5. Atmospheric versus riverine inputs of nitrogen in the Mediterranean

The DIN atmospheric input to the NWMED is in the same range as the Rhone input, which is the major water and nutrient riverine output to the WMED. If the entire WMED is considered (Loÿe-Pilot et al., 1990b), then atmospheric input — derived from field data — is probably higher than that of rivers. Atmospheric input of N ( $1084 \times 10^3 \text{ tons yr}^{-1}$ ) is equal to the riverine input ( $1000 \times 10^3 \text{ tons yr}^{-1}$ ) for the whole Mediterranean according to UNEP/WMO estimations (Bashkin, Erdman, Abramychyev, Sofiev, Pripulina & Gusev, 1997), the atmospheric input being predominant in the southern zones. If the fraction of riverine input which is derived from the atmosphere is taken into account, the atmospheric input is ~60% of the total continental supply of nitrogen to the Mediterranean (Bashkin et al., 1997).

Because riverine nutrients are removed by biological activity in estuarine and proximal coastal zones, atmospheric input is probably the main source of nitrogen for the open sea, even in the northern zones of the Mediterranean.

#### 4.1.6. Impact of atmospheric input of nitrogen on primary production

Atmospheric input of N contributes to new production (Dugdale & Goering, 1967; Eppley & Peterson, 1979; Duce, 1986), which is the primary production fuelled by nutrients originating outside the photic zone, as opposed to production fuelled by recycled nutrients from within the photic zone. Assuming that the assimilation of

nitrogen is in the Redfield ratio to carbon (16/106), then the atmospheric input of nitrogen may account for new production of about  $3.8 \text{ g C m}^{-2} \text{ y}^{-1}$  in the NWMED,  $2.5 \text{ g C m}^{-2} \text{ y}^{-1}$  in the SWMED, and  $2.1 \text{ g C m}^{-2} \text{ y}^{-1}$  in the SEMED.

For coastal zones such as the Gulf of Lions, where total primary production is as high as  $130\text{--}180 \text{ g C m}^{-2} \text{ y}^{-1}$  (Morel, Bricaud, André & Pelaez-Hudlet, 1990; Tusseau & Mouchel, 1995) and new production is approximately one-third of the total, then the atmospheric input of nitrogen represents only about 6–10% of new production, while the Rhône river provides ~50%. However, in oligotrophic zones in the WMED, primary production is about  $50 \text{ g C m}^{-2} \text{ y}^{-1}$  and new production is as low as  $5 \text{ g C m}^{-2} \text{ yr}^{-1}$ . In this case, the atmospheric N input of  $2.5\text{--}3 \text{ g C m}^{-2} \text{ y}^{-1}$  represents up to 60% of new production. This contribution is somewhat lower in the oligotrophic zones of the EMED, where new production has recently been re-estimated at  $16.7 \text{ g C m}^{-2} \text{ y}^{-1}$  (Krom et al., 1991). In this case, the atmospheric input of nitrogen,  $2.1 \text{ g C m}^{-2} \text{ y}^{-1}$  represents ~12% new production.

#### 4.2. Atmospheric input of phosphorus

The atmospheric input of phosphorus has received less attention than that of N, as it is assumed to be less important. The atmospheric input of dissolved inorganic phosphorus to Cap Ferrat is estimated at  $1.4 \mu\text{mol m}^{-2} \text{ d}^{-1}$  and to be between 0.4 and  $1.4 \mu\text{mol m}^{-2} \text{ d}^{-1}$  for the NWMED (Migon, Copin-Montegut, Elégant & Morelli, 1989). Total inorganic phosphorus deposition to the WMED was determined to be  $3.5 \mu\text{mol m}^{-2} \text{ d}^{-1}$  from 3 years of samples collected in Corsica. Crustal material made up 40% of this flux, and the dissolved fraction was estimated to be between one- and two-thirds of the total (Bergametti, Remoudaki, Losno, Steiner, Chatenet & Buat-Ménard, 1992).

The wet deposition of total inorganic phosphorus to the SEMED, estimated from 2 years of measurements on the northern coast of Israel was  $0.6 \mu\text{mol m}^{-2} \text{ d}^{-1}$  and dry deposition was  $1.2 \mu\text{mol m}^{-2} \text{ d}^{-1}$  (Herut & Krom, 1996), giving a total rate of  $1.8 \mu\text{mol m}^{-2} \text{ d}^{-1}$ .

##### 4.2.1. Saharan dust and phosphorus

Phosphorus values of Saharan dust range from 0.04% found in the SEMED to 0.07% from Libyan desert material (Herut & Krom, 1996). Values as high as 0.25–0.40% were found in Saharan dust collected in the NWMED (Loÿe-Pilot, personal communication). A mean content of 0.25% gives a mean phosphorus deposition from Saharan dust in this region of  $2.8 \mu\text{mol m}^{-2} \text{ d}^{-1}$ . However, the question of the solubility and bioavailability of this phosphorus is still open. According to Krom et al. (1991), Saharan dust may act as a trap for soluble phosphorus by adsorption of  $\text{PO}_4^{3-}$  on to Fe-rich particles, whereas Lepple (1971) found that up to 8% of Saharan dust phosphorus is soluble in seawater. From a partitioning study between crustal and anthropogenic phosphorus, Bergametti et al. (1992) estimated that between  $1.2$  and  $2.7 \mu\text{mol P m}^{-2} \text{ d}^{-1}$  is dissolved in seawater from atmospheric input and is available for biota. Similar values ( $1.2 \mu\text{mol P m}^{-2} \text{ d}^{-1}$ ) were obtained from calculations for the EMED, assuming a dust flux of  $36 \text{ g m}^{-2} \text{ yr}^{-1}$ , with 0.4% P and a solubility in seawater of 10% (see Section 5.1).

Table 15

Dissolved inorganic phosphorus (DIP) and total inorganic phosphorus (TIP) deposition (values in  $\mu\text{mol m}^{-2} \text{d}^{-1}$ )

	DIP	TIP
Cap Ferrat <sup>a</sup>	1.4	–
NWMED <sup>b</sup>	0.4–1.4	–
Corsica <sup>c</sup>	1.2–2.7	3.5 (2.1 anthro+1.4 Sahara)
Corsica <sup>d</sup>	–	2.8 Sahara
avg. WMED	1.5	3 (0.035 gP $\text{m}^{-2} \text{yr}^{-1}$ )
EMED <sup>e</sup>	0.6	1.8 (0.021 gP $\text{m}^{-2} \text{yr}^{-1}$ )

<sup>a</sup> Migon (1989).

<sup>b</sup> Migon (1989).

<sup>c</sup> Bergametti et al. (1992).

<sup>d</sup> Loÿe-Pilot, unpublished results.

<sup>e</sup> Herut and Krum (1996).

Table 15 shows estimates of the dissolved inorganic phosphorus (DIP) inputs to the WMED ( $1.5 \mu\text{M P m}^{-2} \text{yr}^{-1}$ ) and EMED ( $0.6 \mu\text{M P m}^{-2} \text{yr}^{-1}$ ), and Table 16 compares total inorganic phosphorus (TIP) atmospheric inputs with riverine P discharge estimates.

#### 4.2.2. Impact of atmospheric phosphorus on primary production

Taking the two values from Bergametti et al. (1992) and applying the Redfield ratio (C/P=106/1) gives potential carbon production as a result of atmospheric dissolved phosphorus input of 0.56 to 1.25 g C  $\text{m}^{-2} \text{yr}^{-1}$ , that is, 11–25% of the new production of the very oligotrophic zones of the WMED, but one to a few percent for the other more productive zones. The impact of the atmosphere therefore seems to be low for phosphorus; certainly lower than for nitrogen.

The high N/P ratio of atmospheric input ( $\sim 70$ ) may be responsible for the high N/P ratio (20–27) observed in Mediterranean seawater (Herut & Krom, 1996).

Table 16

Comparison of riverine (R) and atmospheric (A) inputs of nitrogen and phosphorus (all in tons  $\cdot 10^3 \text{yr}^{-1}$ ). Data on N and P riverine inputs from Vollenweider, Rinaldi, Viviani and Todini (1996); on N atmospheric input from Erdman et al. (1994); on P atmospheric input: from average TIP fluxes in three sub-basins (west to east) of 0.035–0.028–0.021 g P  $\text{m}^{-2} \text{yr}^{-1}$ , respectively

Sub-basin	Area ( $10^6 \text{km}^2$ )	Nitrogen		Phosphorus	
		R	A	R	A
WMED	0.84	577	459	78	29
CMED	0.80	212	346	29	22
EMED	0.70	223	263	30	15
Total	2.34	1011	1068	137	66
% atmosphere			51		33

#### 4.3. Conclusions and hypotheses

The relative importance of atmospheric input of nutrients to the Mediterranean increases from the shoreline to the open sea with the decreasing influence of riverine inputs, and from active mixing zones (margin ‘upwellings’, frontal zones) to oligotrophic ‘stable’ zones. On an annual scale, the nitrogen atmospheric input is significant for oligotrophic zones, in which it may account for up to 60% of new production, whereas atmospheric phosphorus may account for a maximum of 25%.

It is evident that the impact of the atmospheric input of nutrients to the open Mediterranean is difficult to detect, and would in any case be negligible during the period of intense primary production because of mixing with deep rich waters. But during the summer and early fall stratification of the Mediterranean, atmospheric inputs represent a non-negligible source of nutrients (up to  $4000 \mu\text{mol N m}^{-2} \text{d}^{-1}$ ), which may trigger small but detectable phytoplankton blooms (Fig. 10), as reported by Martin et al. (1989).

In this respect, annual atmospheric N budgets may have little significance, but the episodic character of the atmospheric supply may be significant over short time-spans. An example of this is the wet atmospheric N deposition pulse which was studied at Villefranche (Migon et al., 1989). In this case, prior to deposition, oligotrophic conditions persisted and the upper 18 m of the water column was stratified by a low salinity layer. After the deposition event, the atmospheric N signature was clearly observable for 21 days. Neglecting the role of phosphorus, this input of 0.23




Fig. 10. Wet atmospheric input of nitrogen and its effect on productivity (from Martin et al., 1989).

$\mu\text{mol N l}^{-1}$  would theoretically have been able to induce new production of  $16 \text{ mg C m}^{-2} \text{ day}^{-1}$ .

In the same way, the sporadic input of Saharan dust may be a source of phosphorus, especially when originating from mixed (anthropogenic/Saharan) rains, in which case phosphorus would be partly desorbed and dissolved from the dust. It has been suggested that some summer algal blooms may be explained by such Saharan dust fallout (see Section 5.1) (Dulac et al., 1996).

The role of Saharan dust still needs careful assessment, as it may be an important source of phosphorus and iron (see below). The processes and magnitude of phosphorus and iron dissolution from desert dust and their biological impact are important topics to be investigated in the oligotrophic Mediterranean. Two examples of this will be described in the next section.

## 5. Potential effects of atmospheric input on surface ocean biogeochemistry: preliminary speculations

It appears that atmospheric inputs of metals and nutrients to the Mediterranean are of similar magnitude to fluvial inputs, although there are several uncertainties which currently prevent us from quantifying the magnitude and chemical effects of these inputs. The next question that should be addressed is whether they are of significance to biogeochemical cycles in the ocean. We will discuss below two examples in which it seems that atmospheric inputs coincide with a biological response.

### 5.1. *Meteosat and CZAS: a case of P-fertilisation?*

Dulac et al. (1996) presented a study combining monitoring of desert aerosol from *Meteosat* and of marine phytoplankton from CZCS (Coastal Zone Colour Scanner) in the WMED during summer 1983. Daily *Meteosat* images in the solar spectrum (visible channel) were used to quantify African dust transport, following the method described in Moulin et al. (1997a). The CZCS multichannel data were used to monitor marine biology activity in the WMED. Estimates of phytoplankton abundances were based on the algorithm of Morel (1988) using ratios of seawater reflectances at  $0.44$  and  $0.55 \mu\text{m}$  after correction for atmospheric effects (André & Morel, 1991).

Dulac et al. (1996) observed a particularly strong desert dust transport event at the end of July 1983 (peak on Julian day 209), which yielded a strong dust flux to Mediterranean waters, with a cumulative input up to  $1.5 \text{ g m}^{-2}$  during the week, as shown in Fig. 11. They also analysed about one hundred CZCS images of the Ligurian Sea during the same period, and averaged chlorophyll estimates in a region of  $100 \times 100 \text{ km}^2$  near the DYFAMED station (corresponding to about 10000 CZCS pixels). Any images in which more than half the CZCS pixels were cloudy or in which the geometric standard deviation of the pixels was larger than 1.5 were rejected. Any images in which the satellite signal was contaminated by high desert dust loads were also eliminated, using the near-infrared channel ( $0.75 \mu\text{m}$ ) of the

CZCS. In all, nearly 90% of the initial images were rejected; the resulting maps of phytoplankton concentrations are shown in Fig. 12. The peak of dust input (equivalent to approximately  $9 \mu\text{mol P m}^{-2}$ ) was followed, two weeks later, by a significant increase in phytoplankton concentrations, with a peak value of about  $0.5 \text{ mg m}^{-3}$  observed on August 16 (day 228). A response time of one week seems reasonable for phytoplankton growth in natural conditions. It is unlikely that the observed increase in phytoplankton concentration was connected with a decrease in summer surface water stratification, since during the whole period the surface wind speed was always lower than  $1 \text{ m sec}^{-1}$ .

Assuming  $1.5 \text{ g m}^{-2}$  of dust, 0.23% of P in the dust (Graham & Duce, 1982) and a solubility of 8% (Lepple, 1971), produces a flux of  $9 \mu\text{MP m}^{-2}$  during this episode. This corresponds to an addition of  $9 \text{ nM l}^{-1}$  to a 1-m deep layer. This seems too small to produce a bloom (even assuming surface water P values as low as 20 nM). However, we do not really know how much P dissolved from the dust, so the flux may have been larger than the calculations suggest. Another point is that an input into a highly stratified layer would be more significant. While observations show a biological response after the dust storm, at this stage we do not know what the causal relationship was.

## 5.2. Wet deposition of dust: the role of Fe speciation in phytoplankton blooms

The *Cemiliana* theory put forward by Saydam (1996) suggests that the wet deposition of reduced iron from desert dusts to surface waters of the Mediterranean, during the day, has the potential to induce phytoplankton blooms. Blooms have been observed several times in the Mediterranean and even in enclosed basins like the Black Sea, and have been linked with prior wet dust deposition events (Saydam & Yilmaz, 1998; Saydam & Polat, 1999).




Fig. 11. Meteosat-derived weekly maps of dust input to Mediterranean between July 27 and August 2 1983. Continuous input of African dust occurred during this period, peaking on July 30 1983. No significant rain was reported during this period. Meteosat daily images allow estimation of dry deposition flux (Dulac et al., 1996) and geographical distribution is also shown. Corresponding dissolved phosphorus input ranges from 5 to  $10 \mu\text{g l}^{-1}$  for NW Mediterranean surface waters.




Fig. 12. CZCS-derived weekly maps of phytoplankton concentrations during summer 1983: a) mid-June: spring blooms ended, and Ligurian and Adriatic Seas remained relatively productive; b) early August: during dust deposition event, whole basin exhibited low phytoplankton concentrations, except for coastal zone of Adriatic Sea; c) mid-August: 2 weeks after dust deposition event, unusually high concentrations were observed in Ligurian and Adriatic Seas.

An example occurred during the summer of 1997 over the Central Mediterranean and off the coast of Libya. AVHRR and Meteosat scenes from the period 1–2 June 1997 (F. Dulac, pers. comm.) indicate the transport and possible wet deposition of dust during midday over a position where a subsequent *Emiliania huxleyi* bloom was later observed (Fig. 14). Synoptic scale meteorological charts also confirm the presence of atmospheric vertical migration and a cold front over the region, supporting the view that wet deposition occurred.

The *Cemiliana* theory necessitates the presence of ‘desert-originated dust’ for the

production of reduced Fe. Laboratory experiments of light-induced dissolution of hematite, goethite and lepidocrosite (Sulzberger, 1993; Sulzberger & Laubscher, 1995) with oxalate as reductant ligand have shown that, at pH=3, only photochemical formation of dissolved Fe(II) occurs with lepidocrosite, but does not result in appreciable amounts of dissolved Fe(II) when hematite or goethite are used. In confirmation, the revised rainwater sampling strategy of Ozsoy (1998) has shown that iron originating from desert sources may be photochemically reduced within cloud droplets to the +2 oxidation state, as suggested by Faust (1994).

The importance of atmospheric sources in supplying iron to the surface of the open ocean is the subject of some debate. Johnson, Gordon and Coale (1997) showed that dissolved iron profiles in the Pacific and Atlantic Oceans have a uniform shape, with a 'nutrient-type' profile. They agree that the major source of iron to the deep sea is eolian transport, but show that integrated (surface to 500 m) concentrations are only poorly correlated with eolian iron deposition flux. However, Boyle (1997) argues that the distribution of iron in deep waters is not well understood, and that the supply of iron to the open ocean is, in fact, predominantly eolian. Evidence of the impact of atmospheric inputs to particulate Fe levels in the Mediterranean may be seen in the distributions presented by Dulac et al. (1996). In that work, a dramatic increase in surface water particulate Fe was seen in the Ligurian Sea after a major input of Saharan dust in July 1988 (Fig. 13).

Despite the clear impact that atmospheric dust may have on particulate and even dissolved Fe loads in the Mediterranean (see section 4.6), it seems unlikely that Fe will limit biological production in this region, since dissolved Fe levels appear to be relatively high (>1 nM) and macronutrient levels are low. The possibility that it is the deposition of reduced Fe which triggers such events also seems unlikely, since photochemical reduction of existing surface dissolved iron during daylight hours, as proposed by Johnson, Coale, Elrod and Tindale (1994), would already ensure a sufficient supply.

In order to evaluate this postulated relationship between atmospheric/oceanographic forcing functions and biological responses, it is important to establish whether the links are causal. This needs to be done through direct sampling of ocean and the atmosphere, during a strong Saharan dust event and the following weeks.

## 6. Conclusions

Average atmospherically deposited annual mass fluxes of dust are estimated at 8–12–35 g m<sup>-2</sup>, in the Western Mediterranean (WMED), Central Mediterranean (CMED) and Eastern Mediterranean (EMED) respectively. These fluxes are one-third those of riverine sources.

- Available satellite estimates agree well with the few offshore trap measurements.
- Atmospheric dust deposition is the most important part of the geochemical cycles for some elements and substances (e.g. Pb, Cd, PAH).
- Saharan dust pulses not only perturb the atmospheric conditions above the Medit-




Fig. 13. Vertical profile of particulate iron in surface waters of Ligurian Sea, before and after a major atmospheric input of Saharan dust on July 7 1988 (from Dulac et al., 1996).

erranean but, following deposition of this material to the sea surface, they may have a significant effect on the concentration levels of some trace elements in the mixed layer.

- The speciation signatures of some trace metals are different in anthropogenic-rich (European) and crust-rich (Saharan) aerosols; in particular, more of the total concentrations of Cu, Pb and Zn are in exchangeable (potentially mobile) association in the anthropogenic-rich aerosols.
- Atmospherically-transported trace metals are transmitted to marine biogeochemical cycles following the deposition of aerosols at the sea surface by either 'wet' or 'dry' deposition. In both deposition modes, the solubility of the trace metals is constrained by the extent to which they are in exchangeable association in the parent aerosols; but, for the 'wet' deposition, solution pH and particulate load both impose additional controls.
- The importance of atmospheric input of nutrients to the Mediterranean increases from the shoreline to the open sea with the decreasing influence of riverine inputs, and from active mixing zones (margin 'upwellings', frontal zones) to oligotrophic 'stable' zones. On an annual scale, the nitrogen atmospheric input is significant for oligotrophic zones, where it may account for up to 60% of new production, whereas atmospheric phosphorus may only account for a maximum of 25%.
- Biological effects linked with wet and dry atmospheric inputs of nutrients are suggested.

### 6.1. Future work

Unique information on the geographic source and tropospheric transport of particulate plumes is provided by satellite observations, and the recently launched ocean




Fig. 14. Four images representing respectively (from top): (a) NOAA-AVHRR composite imagery (June 9 1997), illustrating the high reflection region; (b) CEA Meteosat quicklook showing the massive dust outbreak over the Sahara and small filaments extending towards the Gulf of Sirte (June 2 1997); (c) dust loading prediction (June 2 1997) from MEDUSE model (<http://halo.hi.is/meduse/index.html>); and (d) possible wet deposition (June 2 1997) over a vast area offshore Libya, as derived from NOAA-AVHRR composite imagery (VIS and IR). Dust outbreaks in satellite image and deposition model was delivered to sea with rain, and may have triggered *Emiliania huxleyi* bloom observed one week later.

and multispectral sensors (OCTS, POLDER, SEAWIFS) will allow more extensive and precise monitoring.

In order to provide insight on the impact of dust events on the chemistry and climatology of an enclosed sea such as the Mediterranean, satellite observations should be:

1. coupled to independent source tracers, including air mass trajectories and chemical tracers such as Sr, Nd and Pb isotopes, and
2. associated with nutrient and trace metal measurements at the onset and during specific events.

Certain specific actions are also recommended (CIESM, 1997):




Fig. 14. (continued)

1. expansion of the present rain/dust collection network in the EMED and SMED regions, to provide more complete regional coverage of dust deposition of the basin;
2. a well-coordinated multidisciplinary research effort should be planned aimed at sampling the atmosphere and the surface Mediterranean in real time during and immediately after a Saharan dust event, to take advantage of this natural seeding experiment.
3. a data base should be established, to file existing atmospheric and oceanic information in the Mediterranean; this data base should be available to all workers in the field and could be supported by an international commission such as CIESM (Commission Internationale pour l'Exploration Scientifique de la mer Méditerranée).

## Acknowledgements

This work was partially supported by MTP-MATER Contract MAS3-CT96-0051, STEP Contract CT N-080-0080, and World Meteorological Organisation Contract 20.028/A/CNS. This is IGM-CNR scientific contribution no. 1121. We also thank G. Rampazzo and S. Cristini for discussions and trace metal data, G. Cesari for sampling help, G. Quarantotto for chemistry work, G. Zini and E. Masetti for maps and drawings, and G. Walton for revision of the English text. We also acknowledge an anonymous reviewer for useful suggestions. Lastly, we thank Frédéric Briand (CIESM director) and Scott Fowler for encouraging us to initiate this work and for supporting it.

## References

- Andersen, V., & Nival, P. (1988). Modèles d'écosystème pélagique des eaux côtières de la mer Ligure. In H. J. Minas, & P. Nival, *Océanographie Pélagique Méditerranéenne*, *Oceanologica Acta*, pp. 211–217.
- André, J.-M., & Morel, A. (1991). Atmospheric corrections and interpretation of marine radiances in CZCS imagery, revisited. *Oceanologica Acta*, *14*, 3–22.
- Arimoto, R., & Duce, R. A. (1986). Dry deposition models and the air-sea exchange of trace elements. *Journal of Geophysical Research*, *88*, 2787–2792.
- Axenfeld, F., Munch, J., Pacyna, J. M., Duiser, J. A., & Veldt, C. (1992). *Emissionsdatenbasis für die Spurenelemente As, Cd, Hg, Pb, Zn und für spezielle organische Verbindungen Lindan, HCB, PCB und PAK*, Forschungsbericht 104 02 588, Umweltforschungsplan des Bundesministers für Umwelt Naturschutz und Reaktorsicherheit, Dornier, Friedrichshafen, 103 pp.
- Baeyens, W., Dehairs, F., & Dedeurwaerder, H. (1990). Wet and dry deposition fluxes above the North Sea. *Atmospheric Environment*, *24A*, 1693–1703.
- Bashkin, V. N., Erdman, L. K., Abramychiev, A. Yu., Sofiev, M. A., Pripulina, I. V., & Gusev, A. V. (1997). The input of anthropogenic airborne nitrogen to the Mediterranean Sea through its watershed. *MAP Technical Reports Series*, Vol. 118, UNEP, Athens (95 pp).
- Bergametti, G. (1987). *Apports de matière par voie atmosphérique à la Méditerranée Occidentale*. Ph.D. thesis, Univ. Paris 7.
- Bergametti, G., Dutot, A. L., Buat-Ménard, P., Losno, R., & Remoudaki, E. (1989). Seasonal variability of the elemental composition of atmospheric aerosol particles over the North-western Mediterranean. *Tellus*, *41B*, 353–361.
- Bergametti, G., Remoudaki, E., Losno, R., Steiner, E., Chatenet, B., & Buat-Ménard, P. (1992). Source, transport and deposition of atmospheric phosphorus over the north-western Mediterranean. *Journal of Atmospheric Chemistry*, *14*, 501–513.
- Berland, B. R., Bonin, D. J., & Maestrini, S. Y. (1980). Azote ou phosphore? Considérations sur le 'paradoxe nutritionnel' de la mer méditerranée. *Oceanologica Acta*, *3*, 135–142.
- Berland, B. R., Bonin, D. J., Maestrini, S. Y., & Pointier, J. P. (1973). Etude de la fertilité des eaux marines au moyen de tests biologiques effectués avec des cultures d'algues. IV. Etude des eaux côtières méditerranéennes. *Internationale Revue des Gesamten Hydrobiologie*, *58*, 473–500.
- Boyle, E. (1997). What controls dissolved iron concentrations in the world ocean? A comment. *Marine Chemistry*, *57*, 163–167.
- Boyle, E. A., Chapnick, S. D., Bai, X. X., & Spivack, A. (1985). Trace metal enrichments in the Mediterranean Sea. *Earth and Planetary Sciences Letters*, *74*, 405–419.
- Bruland, K. W., Donat, J. R., & Hutchins, D. A. (1991). Interactive influences of bioactive trace metals on biological production in oceanic waters. *Limnology and Oceanography*, *36*, 1555–1577.
- Buat-Ménard, P. (1993). Global change in atmospheric metal cycles. In C. N. Hewitt, & W. T. Sturges, *Global atmospheric chemical change* (pp. 271–309).

- Caquineau, S., Gaudichet, A., Gomes, L., Magonthier, M.-C., & Chatenet, B. (1999). Saharan dust, Part 1: clay ratio as a relevant tracer to assess the origin of soil-derived aerosols. *Geophysical Research and Letters*, (in press).
- Carratala, A., Bellot, J., Gomez, A., & Millan, M. (1996). African dust influence on rainwater on the eastern coast of Spain. In S. Guerzoni, & R. Chester, *The impact of desert dust across the Mediterranean* (pp. 323–332). Kluwer Academic Publishers.
- Chester, R., Nimmo, M., & Murphy, K. J. T. (1989). Atmospheric inputs of trace metals, trace organics and nutrients to the western Mediterranean — The EROS 2000 Programme. *Water Pollution Research Reports*, 13, 357–367.
- Chester, R., Nimmo, M., & Keyse, S. (1996). The influence of Saharan and eastern desert-dust on the trace metal composition of Mediterranean aerosols and rainwater: an overview. In S. Guerzoni, & R. Chester, *The Impact of Desert Dust Across the Mediterranean* (pp. 253–273). Kluwer Academic Publishers.
- Chester, R., Sharples, E. J., Sanders, G. S., & Saydam, A. C. (1984). Saharan dust incursion over the Tyrrhenian Sea. *Atmospheric Environment*, 18, 929–935.
- Chester, R., Nimmo, M., Murphy, K. J. T., & Nicolas, E. (1990). Atmospheric trace metals transported to the western Mediterranean: data from a station on Cap Ferrat. *Water Pollution Research Reports*, 20, 597–612.
- Chester, R., Murphy, K. J. T., Lin, F. J., Berry, A. S., Bradshaw, G. A., & Corcoran, P. A. (1993). Factors controlling solubilities of trace metals from non-remote aerosols deposited to the sea surface by the 'dry' deposition mode. *Marine Chemistry*, 42, 107–126.
- Chester, R., Baxter, G. G., Behairy, A. K. A., Connor, K., Cross, D., Elderfield, H., & Padgham, R. C. (1977). Soil-sized eolian dusts from the lower troposphere of the Eastern Mediterranean Sea. *Marine Geology*, 24, 201–217.
- Chou, L., & Wollast, R. (1997). Biogeochemical behaviour and mass balance of dissolved aluminium in the western Mediterranean. *Deep-Sea Research II*, 44, 741–768.
- CIESM (1997). Role of ocean-atmosphere processes in the biogeochemistry of the Mediterranean Sea, *CIESM Workshop Series*, Vol. 3, (29 pp).
- Coale, K. H., Johnson, K. S., Fitzwater, S. E., Gordon, R. M., Tanner, S., Chavez, F. P., Ferioli, L., Sakamoto, C., Rogers, P., Millero, F., Steinberg, P., Nightingale, P., Cooper, D., Cochlan, W. P., Landry, M. R., Constantinou, J., Rollwagen, G., Trasvina, A., & Kudela, R. (1996). A massive phytoplankton bloom induced by an ecosystem-scale iron fertilization experiment in the equatorial Pacific Ocean. *Nature, London*, 383, 495–501.
- Cornell, S., Rendell, A., & Jickells, T. (1995). Atmospheric inputs of dissolved organic nitrogen to the oceans. *Nature, London*, 376, 243–246.
- Dolan, J. R., Thingstad, T. F., & Rassoulzadegan, F. (1995). Phosphate transfer between size-fractions in Villefranche Bay (N W Mediterranean Sea), France in autumn 1992. *Ophelia*, 41, 71–85.
- Dolske, D. A., & Gatz, D. F. (1985). A field intercomparison of methods for the measurement of particle and gas dry deposition. *Journal of Geophysical Research*, 90, 2076–2084.
- Dorten, W., Elbaz-Poulichet, F., Mart, L., & Martin, J. M. (1991). Reassessment of the river input of trace metals into the Mediterranean Sea. *Ambio*, 20, 2–6.
- Duce, R. A. (1986). In P. Buat-Ménard, *The impact of atmospheric nitrogen, phosphorus and iron species on marine biological productivity* (pp. 497–529). *The role of air-sea exchange in geochemical cycling, NATO ASI Series, Serie C*, 185.
- Duce, R.A. (1997). *Atmospheric input of pollution to the oceans*. Meeting of the World Meteorological Organization Commission for Marine Meteorology, Havana, Cuba, 17 March 1997.
- Duce, R. A., Unni, C. K., Ray, B. J., & Arimoto, R. (1982). Atmospheric deposition of trace metals to the Tropical North Pacific. *Eos*, 63, 1987.
- Duce, R. A., Liss, P. S., Merrill, J. T., Atlas, E. L., Buat-Ménard, P., Hicks, B. B., Miller, J. M., Prospero, J. M., Arimoto, R., Church, T. M., Ellis, W., Galloway, J. N., Hansen, L., Jickells, T. D., Knap, A. H., Reinhardt, K. H., Schneider, B., Soudine, A., Tokos, J. J., Tsunogai, S., Wollast, R., & Zhou, M. (1991). The atmospheric input of trace species to the world ocean. *Global Biogeochemical Cycles*, 5, 193–259.

- Dugdale, R. C., & Goering, J. J. (1967). Uptake of new and regenerated forms of nitrogen in primary productivity. *Limnology and Oceanography*, 9, 170–184.
- Dulac, F., Buat-Ménard, P., Arnold, M., & Ezat, U. (1987). Atmospheric input of trace metals to the Western Mediterranean Sea: 1. Factors controlling the variability of atmospheric concentrations. *Journal of Geophysical Research*, 92, 8437–8453.
- Dulac, F., Buat-Ménard, P., Ezat, U., & Bergametti, G. (1989). Atmospheric input of trace metals to the Western Mediterranean Sea: uncertainties in modelling dry deposition from cascade impactor data. *Tellus*, 41B, 362–378.
- Dulac, F., Moulin, C., Lambert, C. E., Guillard, F., Poitou, J., Guelle, W., Quénel, C., Schneider, X., & Ezat, U. (1996). Quantitative remote sensing of African dust transport to the Mediterranean. In S. Guerzoni, & R. Chester, *The impact of desert dust across the Mediterranean* (pp. 25–49). Kluwer Academic Publishers.
- Elbaz-Poulichet, F., Guan, D. M., Seyler, P., Martin, J. M., Morley, N., Statham, P. J., Burton, J. D., Mart, L., & Klahre, P. (1989). Dissolved trace metals and metalloids in the Rhône River/estuarine system. *Water Pollution Research Reports*, 13, 357–367.
- EMEP (1994) *EMEP Workshop on the Accuracy of Measurements*, Passau, Germany, November 1993, EMEP/CCC Report 2/94.
- Eppley, R. W., & Peterson, B. J. (1979). Particulate organic matter flux and planktonic new production. *Nature, London*, 282, 677–680.
- Erdman, L. K., Sofiev, M., Subbotin, S., Dedkova, I., Afinogenova, O., Chesukina, T., Pavlovskaya, L., & Soudine, A. (1994). Assessment of Airborne Pollution of the Mediterranean Sea by Sulphur and Nitrogen Compounds and heavy metals in 1991. *MAP Technical Report Series*, UNEP/WMO, Vol. 85, Athens (304 pp).
- Faust, B. (1994). Photochemistry of clouds, fogs, and aerosols. *Environmental Science and Technology*, 28, 217–222.
- Fiala, M., Cahet, G., Jacques, G., Neveux, J., & Panouse, M. (1976). Fertilisation de communautés phytoplanctoniques. 1. Cas d'un milieu oligotrophe: Méditerranée nord-occidentale. *Journal of Experimental Marine Biology and Ecology*, 24, 151–163.
- Fisher, D., Ceraso, J., Mathew, T., & Oppenheimer, M. (1988). *Polluted coastal waters: the role of acid rain*. New York: Environmental Defense Fund.
- Ganor, E., & Foner, H. (1996). The mineralogical and chemical properties and the behaviour of the aeolian Saharan Dust over Israel. In S. Guerzoni, & R. Chester, *The impact of desert dust across the Mediterranean* (pp. 173–182). Kluwer Academic Publishers.
- GESAMP (1989). The atmospheric input of trace species to the world ocean, *WMO Reports and Studies*, (Vol. 38), (111 pp).
- Graham, W. F., & Duce, R. A. (1982). The atmospheric transport of phosphorus to the western North Atlantic. *Atmospheric Environment*, 16, 1089–1097.
- Guerzoni, S., Lenaz, R., & Quarantotto, G. (1988). Field measurements at sea: atmospheric trace metals 'end-members' in the Mediterranean. *Air Pollution Research Report*, 14, 96–100.
- Guerzoni, S., Molinaroli, E., & Chester, R. (1997). Saharan dust inputs to the western Mediterranean Sea: depositional patterns, geochemistry and sedimentological implications. *Deep-Sea Research II*, 44, 631–654.
- Guerzoni, S., Quarantotto, G., Cesari, G., Molinaroli, E., Rampazzo, G., & Le Bolloch, O. (1996). Trace metal composition and grain-size of particulates in aerosols and precipitation collected in N.W. Mediterranean (39°N, 9°E). In S. Guerzoni, & R. Chester, *The impact of desert dust across the Mediterranean* (pp. 333–338). Kluwer Academic Publishers.
- Guerzoni, S., Molinaroli, E., Rossini, P., Rampazzo, G., Quarantotto, G., & Cristini, S. (1999). Role of desert aerosol in metal fluxes in the Mediterranean area. *Chemosphere*, 39, 229–246.
- Guerzoni, S., Landuzzi, W., Lenaz, R., Quarantotto, G., Rampazzo, G., Molinaroli, E., Turetta, C., Visin, F., Cesari, G., & Cristini, S. (1993). Fluxes of soluble and insoluble metals and nutrients from the atmosphere to the Central Mediterranean Sea. *Water Pollution Research Reports*, 30, 253–260.
- Guieu, C. (1991). *Apports atmosphériques à la Méditerranée nord-occidentale*, Ph.D. thesis, Univ. Paris 6.
- Guieu, C., Thomas, A. J., Martin, J. M., & Brun-Cottan, J. C. (1991). Multielemental characterization of

- the atmospheric input to the Gulf of Lions. Impact upon the upper layer water chemistry. *Water Pollution Research Reports*, 28, 505–518.
- Guieu, C., Chester, R., Nimmo, M., Martin, J.-M., Guerzoni, S., Nicolas, E., Mateu, J., & Keyse, S. (1997). Atmospheric input of dissolved and particulate metals to the north-western Mediterranean. *Deep-Sea Research II*, 44, 655–674.
- Herut, B., & Krom, M. (1996). Atmospheric input of nutrient and dust to the SE Mediterranean. In S. Guerzoni, & R. Chester, *The impact of desert dust across the Mediterranean* (pp. 349–358). Kluwer Academic Publishers.
- Heussner, S., & Monaco, A. (1996). The MTP sediment trap experiments: highlights on particle fluxes through the eutrophic and oligotrophic systems of the Mediterranean Sea. In P. Wassmann, & T. Tselepidis, *Second Workshop of MTP-MAST Programme — Extended abstracts*, pp. 233–239.
- Hydes, D. J., De Lange, G. J., & Debaar, H. J. W. (1988). Dissolved aluminium in the Mediterranean. *Geochimica et Cosmochimica Acta*, 52, 2107–2114.
- Johnson, K. S., Coale, K. H., Elrod, V. A., & Tindale, N. W. (1994). Iron photochemistry in seawater from the equatorial Pacific. *Marine Chemistry*, 46, 319–334.
- Johnson, K. S., Gordon, R. M., & Coale, K. H. (1997). What controls dissolved iron concentrations in the world ocean? *Marine Chemistry*, 57, 137–161.
- Keyse, S. (1995) *Trace metal chemistry of Mediterranean rain waters*, Ph.D. thesis, Univ. of Liverpool, UK.
- Klein, C. (1998) *Apports atmosphériques en azote inorganique dissous: dépôt sec par les aérosols et effet des précipitations sur le réseau microbien*, Thèse Université P. et M. Curie, Paris.
- Krom, M. D., Kress, N., Brenner, S., & Gordon, L. I. (1991). Phosphorus limitation of primary productivity in the eastern Mediterranean Sea. *Limnology and Oceanography*, 36, 424–432.
- Kubilya, N. S., & Saydam, A. C. (1995). Trace elements in atmospheric particulates over the Eastern Mediterranean; concentrations, sources, and temporal variability. *Atmospheric Environment*, 29, 2289–2300.
- Kumar, N., Anderson, R. F., Mortlock, R. A., Froelich, P. N., Kubik, P., Ditttrich-Hannen, B., & Suter, M. (1995). Increased biological productivity and export production in the glacial Southern Ocean. *Nature, London*, 378, 675–680.
- Le Bolloch, O., & Guerzoni, S. (1995). Acid and alkaline deposition in precipitation on the western coast of Sardinia, Central Mediterranean (40°N, 8°E). *Water Air and Soil Pollution*, 85, 2155–2160.
- Lepple, F. K. (1971). *Eolian dust over the North Atlantic Ocean*. Ph.D. thesis, Univ. of Delaware.
- Lim, B., Jickells, T. D., Colin, J. L., & Losno, R. (1994). Solubilities of Al, Pb, Cu, and Zn in rain sampled in the marine environment over the North Atlantic Ocean and Mediterranean Sea. *Global Biogeochemical Cycles*, 8, 349–362.
- Loÿe-Pilot, M. D., & Martin, J. M. (1996). Saharan dust input to the western Mediterranean: an eleven-year record in Corsica. In S. Guerzoni, & R. Chester, *The impact of desert dust across the Mediterranean* (pp. 191–199). Kluwer Academic Publishers.
- Loÿe-Pilot, M. D., Martin, J. M., & Morelli, J. (1986). Influence of Saharan dust on the rain acidity and atmospheric input to the Mediterranean. *Nature, London*, 321, 427–428.
- Loÿe-Pilot, M. D., Martin, J. M., & Morelli, J. (1990a). Atmospheric wet deposition of inorganic nitrogen to the North Western Mediterranean Basin. *Water Pollution Research Reports*, 20, 623–634.
- Loÿe-Pilot, M. D., Martin, J. M., & Morelli, J. (1990b). Atmospheric input of inorganic nitrogen to the Western Mediterranean. *Biogeochemistry*, 9, 117–134.
- Loÿe-Pilot, M. D., Cauwet, G., Spitz, A., & Martin, J. M. (1992). Preliminary results on atmospheric wet deposition of organic carbon and nitrogen in Corsica. *Water Pollution Research Report*, 28, 519–532.
- Mallin, M. A., Paerl, H. W., Rudek, J., & Bates, P. W. (1993). Regulation of estuarine primary production by watershed rainfall and river flow. *Marine Ecology Progress Series*, 93, 199–203.
- Martin, J. H. (1990). Glacial-interglacial CO<sub>2</sub> change: the iron hypothesis. *Paleoceanography*, 5, 1–13.
- Martin, J. M., Elbaz-Poulichet, F., Guieu, C., Loÿe-Pilot, M. D., & Han, G. (1989). River versus atmospheric input of material to the Mediterranean: an overview. *Marine Chemistry*, 28, 159–182.
- Mateu, J., Colom, M., Forteza, R., & Cerda, V. (1993). Monitoring of major and minor inorganic components of aerosols from the Mallorca station. *Water Pollution Research Reports*, 30, 261–270.
- Mazzucotelli, A., Landuzzi, W., Lenaz, R., Oliveri, F., Tomadin, L., & Vannucci, E. R. (1986). Polveri

- in sospensione nella bassa atmosfera del Mare Tirreno e del Canale di Sicilia (crociera Ban 80). *Memorie della Società Geologica Italiana*, 27, 311–321.
- Measures, C. I., & Edmond, J. M. (1988). Aluminium as a tracer of the deep outflow from the Mediterranean. *Journal of Geophysical Research*, 93, 591–595.
- Medinets, V. I. (1996). Shipboard derived concentrations of sulphur and nitrogen compounds and trace metals in the Mediterranean aerosol. In S. Guerzoni, & R. Chester, *The impact of desert dust across the Mediterranean* (pp. 369–373). Kluwer Academic Publishers.
- Migon, C. (1993). Riverine and atmospheric inputs of heavy metals to the Ligurian Sea. *The Science of the Total Environment*, 138, 289–299.
- Migon, C., & Caccia, J. L. (1990). Separation of anthropogenic and natural emissions of particulate heavy metals in the western Mediterranean atmosphere. *Atmospheric Environment*, 24A, 399–405.
- Migon, C., & Nicolas, E. (1998). Effects of antipollution policy on anthropogenic lead transfers in the Ligurian Sea. *Marine Pollution Bulletin*, 36, 775–779.
- Migon, C., Journel, B., & Nicolas, E. (1997). Measurement of trace metal wet, dry and total atmospheric fluxes over the Ligurian Sea. *Atmospheric Environment*, 31, 889–896.
- Migon, C., Copin-Montegut, G., Elégant, L., & Morelli, J. (1989). Etude de l'apport atmosphérique en sels nutritifs au milieu côtier méditerranéen et implications biogéochimiques. *Oceanologica Acta*, 12 (2), 187–191.
- Migon, C., Morelli, J., Nicolas, E., & Copin-Montégut, G. (1991). Evaluation of total atmospheric deposition of Pb, Cd, Cu and Zn to the Ligurian Sea. *The Science of the Total Environment*, 105, 135–148.
- Migon, C., Alleman, L., Leblond, N., & Nicolas, E. (1993). Evolution of atmospheric lead over the NW Med between 1986 and 1992. *Atmospheric Environment*, 27A (14), 2161–2167.
- Mihalopoulos, N., Stephanou, E., Kanakidou, M., Pilitsidis, S., & Bousquet, P. (1997). Tropospheric aerosol ionic composition in the Eastern Mediterranean region. *Tellus*, 49B, 314–326.
- Milliman, J. D., & Martin, J. M. (1997). Topical studies in Oceanography. *Deep-Sea Research II*, 44, 521–950.
- Milliman, J.D., Rutkowsky, C.M., & Meybeck, M. (1995). River discharge to the sea. A global river index (GLORI). *LOICZ Reports and Studies*, (Vol. 4), (pp. 132).
- Molinarioli, E. (1996). Mineralogical characterisation of Saharan dust with a view to its final destination in Mediterranean sediments. In S. Guerzoni, & R. Chester, *The impact of desert dust across the Mediterranean* (pp. 153–162). Kluwer Academic Publishers.
- Molinarioli, E., & Ibbá, A. (1995). Occurrence of palygorskite in aerosol dust (dry and wet) of desert source in SE Sardinia, W Mediterranean. *Giornale di Geologia*, 57, 67–76.
- Morel, A. (1988). Optical modelling of the upper ocean in relation to its biogenous matter content (Case 1 waters). *Journal of Geophysical Research*, 93, 10747–10768.
- Morel, A., Bricaud, A., André, J. M., & Pelaez-Hudlet, J. (1990). Spatial/temporal evolution of the Rhone plume as seen by CZCS imagery — Consequences upon the primary production in the Gulf of Lions. *Water Pollution Research Reports*, 20, 45–62.
- Morley, N. H., Burton, J. D., Tankere, S. P. C., & Martin, J.-M. (1997). Distribution and behaviour of some dissolved trace metals in the western Mediterranean Sea. *Deep-Sea Research II*, 44, 675–691.
- Moulin, C., Guillard, F., Dulac, F., & Lambert, C. E. (1997a). Long-term daily monitoring of Saharan dust load over ocean using Meteosat ISCCP-B2 data, 1. Methodology and preliminary results for 1983–1994 in the Mediterranean. *Journal of Geophysical Research*, 102, 16947–16958.
- Moulin, C., Lambert, E. C., Dulac, F., & Dayan, U. (1997b). Control of atmospheric export of dust from North Africa by the North Atlantic Oscillation. *Nature, London*, 387, 691–694.
- Nicolas, E., Migon, C., Leblond, N., & Journel, B. (1995). Seasonality of dry and wet depositions of trace metals in the Ligurian Sea. *Water Pollution Research Reports*, 32, 275–285.
- Nihlen, T., & Mattsson, J. O. (1989). Studies on eolian dust in Greece. *Geografiska Annaler*, 71, 269–274.
- Nriagu, J. O., & Pacyna, J. M. (1988). Quantitative assessment of worldwide contamination of air, water and soils by trace metals. *Nature, London*, 333, 134–139.
- Owens, N. J. P., Rees, A. P., Woodward, E. M. S., & Mantoura, R. F. C. (1989). Size-fractionated primary production and nitrogen assimilation in the Northwest Mediterranean Sea during January 1989. *Water Pollution Research Bulletin*, 13, 126–135.
- Paerl, H. W., Fogel, M. L., & Bates, P. W. (1993). Atmospheric nitrogen deposition in coastal waters:

- implications for marine primary production and C flux. In R. Guerrero, & C. Pedros-Alio, *Trends in microbial ecology* (pp. 459–464). Spanish Society for Microbiology.
- Patterson, C. C., & Settle, D. M. (1987). Review of data on eolian fluxes of industrial and natural lead to the lands and seas in remote region on a global scale. *Marine Chemistry*, 22, 137–162.
- Pye, K. (1992). Aeolian dust transport and deposition over Crete and adjacent parts of the Mediterranean Sea. *Earth Surface Processes and Landforms*, 17, 271–288.
- Raimbault, P., & Coste, B. (1990). Very high values of the nitrate to phosphorus ratio (>30) in the subsurface layers of the western Mediterranean Sea. *Rapports et procès-verbaux de la réunion de la Commission Internationale d'Exploration Scientifique de la mer Méditerranée*, (Vol. 32), (p. C18).
- Remoudaki, E. (1990). *Etude des processus contrôlant la variabilité temporelle des fluxes atmosphériques de polluants et de poussières minérales en Méditerranée occidentale*, Ph.D. thesis, Univ. Paris 7.
- Roda, F., Bellot, J., Avila, A., Escarré, A., Pinol, J., & Terradas, J. (1993). Saharan dust and the atmospheric inputs of elements and alkalinity to Mediterranean ecosystems. *Water, Air and Soil Pollution*, 66, 277–288.
- Rutten, A., De Lange, G. J., Ziveri, P., Thomson, J., Van Santvoort, P., Colley, S., & Corselli, C. (1999). Recent terrestrial and carbonate fluxes in the pelagic eastern Mediterranean; a comparison between sediment trap and bottom sediment. *Palaeogeography Palaeoclimatology Palaeoecology*, (in press).
- Sandroni, V., & Migon, C. (1997). Significance of trace metal medium-range transport in the western Mediterranean. *The Science of the Total Environment*, 196, 83–89.
- Saydam, A. C. (1996). Can we predict harmful algae blooms. *Harmful Algae News*, 15, 5–6.
- Saydam, A. C., & Yilmaz, A. (1998). *Dust induced algae growth and the Black Sea*. *International Symposium on Fisheries Ecology*. Trabzon, Turkey, 2–4 September 1998.
- Saydam, A. C., & Polat, I. (1999). The impact of saharan dust on the occurrence of algae blooms. In P. M. Borrell, & P. Borrell, *Proceedings of EUROTRAC Symposium '98* (pp. 656–664). Southampton: WIT press.
- Sharp, W. E., & Nardi, G. (1987). A study of the heavy metal pollution in the bottom sediments at Porto di Bagnoli (Naples), Italy. *Journal of Geochemical Exploration*, 29, 31–48.
- Slinn, W. G. N. (1983). *Air to sea transfer of particles*. NATO ASI series, (Vol. 108), (pp. 299–396), Reidel, Dordrecht.
- Spitz, A., Ludwig, W., & Wobig, M. (1990). Survey of amino acid nitrogen in water, sediment and aerosol from the Gulf of Lions. *Water Pollution Research Report*, 20, 187–193.
- Spokes, L. J., & Jickells, T. D. (1996). Factors controlling the solubility of aerosol trace metals in the atmosphere and on mixing into seawater. *Aquatic Geochemistry*, 1 (4), 355–374.
- Sulzberger, B. (1993). The role of heterogeneous photochemical reactions for the formation of dissolved iron species in aquatic systems. *Water Pollution Research Reports*, 30, 183–186.
- Sulzberger, B., & Laubscher, H. (1995). Reactivity of various types of iron(III) (hydr)oxides towards light-induced dissolution. *Marine Chemistry*, 50, 103–115.
- Taylor, S. R. (1964). Abundance of chemical elements in the continental crust: A new table. *Geochimica and Cosmochimica Acta*, 28, 1273–1285.
- Thingstad, T. F., & Rassulzadegan, F. (1995). Nutrient limitations, microbial foodwebs and 'biological pumps': suggested interactions in a P-limited Mediterranean. *Marine Ecology Progress Series*, 117, 299–306.
- Tomadin, L. (1981). Provenance and dispersal of clay minerals in recent sediments of the Central Mediterranean Sea. In F. C. Wezel, *Sedimentary basins of Mediterranean margins*, pp. 313–324.
- Tomadin, L., & Borghini, M. (1987). Source and dispersal of clay minerals from present and Late Quaternary sediments of Southern Adriatic Sea. In: *6th Meeting of the European Clay Groups, Spain*, pp. 537–538.
- Tomadin, L., & Lenaz, R. (1989). Eolian dusts over the Mediterranean and their contribution to the present sedimentation. In M. Leinen, & M. Sarnthein, (pp. 267–282). *Modern and past patterns of global atmospheric transport*, NATO ASI Series, Series C, 282. Kluwer Academic Publishers.
- Tusseau, M. H., & Mouchel, J. M. (1995). Nitrogen inputs to the Gulf of Lions via the Rhone river. *Water Pollution Research Reports*, 3, 49–59.
- UNEP (1984). Pollutants from land-based sources in the Mediterranean. *UNEP Regional Seas Reports and Studies* No. 32.

- Venkatarathnam, K., & Ryan, W. B. F. (1971). Dispersal patterns of clay minerals in the sediments of the Eastern Mediterranean Sea. *Marine Geology*, 11, 261–282.
- Vollenweider, R. A., Rinaldi, A., Viviani, R., & Todini, E. (1996). Assessment of the state of eutrophication in the Mediterranean Sea. *Mediterranean Action Plan Technical Reports Series*, No. 106.
- Windom, H., Smith, R. Jr., Rawlingson, C., Hungspreugs, M., Dharmvanij, S. & Wattayakorn, G. (1988). Trace metal transport in a tropical estuary. *Marine Chemistry*, 24, 293–305.
- Woodward, E. M. S. & Owens, N. J. P. (1989). The influence of the river Rhone upon the nutrient fluxes of the Golfe du Lion. *Water Pollution Research Bulletin*, 13, 79–86.
- Yaaqub, R. R., Davies, T. D., Jickells, T. D., & Miller, J. M. (1991). Trace elements in daily collected aerosols at a site in south east England. *Atmospheric Environment*, A25, 985–996.
- Ziveri, P., De Lange, G. J., Rossignol-Strick, M., Schrader, H., & Thomson, J. (1996). Biogeochemical cycles in the eastern Mediterranean; Actuo (seasonal/annual) variations. 2nd Workshop of Mediterranean Target Project, pp. 217–221.