

UDK: 597 : 639.2 (563.22)
Original scientific paper

**APPLICATION OF MAHALANOBIS DISTANCE FUNCTION
FOR THE MORPHOMETRIC SEPARATION OF SILVERBELLY
(LEIOGNATHUS KLUNZINGERI STEINDACHNER) STOCKS
IN THE GULF OF MERSIN**

PRIMJENA MAHALANOBIS-OVE FUNKCIJE DISKRIMINACIJE
NA ODREĐIVANJE MORFOMETRIJSKIH RAZLIKA IZMEĐU STOKOVA
LEIOGNATHUS KLUNZINGERI STEINDACHNER U ZALJEVU MERSIN

Dursun Avsar, Ferit Bingel and Mustafa Ünsal
Institute of Marine Sciences, M. E. T. U., Erdemili İçel, Turkey

Stock differentiation studies of silverbelly from four localities in the northeastern Mediterranean has been carried out morphometrically applying the Generalized Distance of Mahalanobis and shown that silverbellies of Mersin Bay consisted of four different stocks.

INTRODUCTION

L. klunzingeri is one of the most common Indo-Pacific demersal species in the northeastern Mediterranean. The mean catch of this species during the years 1980—1982 period constituted 10.2% in number of the individuals in Goksu, 9.8% in Tirtar, 3.0% in Seyhan and 3.0% in Tuzla (Bingel, 1987). However, because of its small size, (50—70 mm) these fish have to date no commercial importance in Turkey although it might have some economic potential in future (Bingel, 1981).

These fish have ripe gonads during the summer (Ben-Tuvia 1966). The eggs are pelagic and the spawning occurs from the beginning of April until the end of May (Aksiray, 1954). However, from the length frequency curve fitting data given by Bingel (1987), one may conclude that these fish spawn eventually in two main periods; in April-May and August-September.

Silverbelly have come from the Red Sea along the Gulf of Suez to the eastern basin of the Mediterranean in 1898 (Steindachner, 1898). However, their first detection in Iskenderun Bay was made by Erazi (1943). This species has adapted to the new habitats and spreaded considerably. *Leiognathus klunzingeri* in the Mediterranean are distributed in the eastern part, from Port Said to Mersin Bay and Aegean Sea (Ben-Tuvia, 1966). In fact, one specimen was collected from the neighbourhood of Lampedusa, which is near

ODIU-DBB METOPHANESI
METU-DMS LIBRARY

the eastern coast of Tunisia. These adaptations to the new oceanographic environment may cause the changes in their morphometric characters as was also pointed out by Kosswig (1974). The aim of this study is to define the stock composition of *Leiognathus klunzingeri* in the northern Cilician Basin (Mersin Bay).

Population dynamics of the local populations of each water product shows great diversity (Ricker, 1975). Therefore, the stocks studied, must be differentiated and the unity of the stock or stocks should be stated. In fact, the identification and delimitation of unit stocks are necessary for the assessment of the exploited fish stocks.

MATERIAL AND METHODS

Sample collections were done from the Bay of Mersin, which is located in the northern Mediterranean. Four sampling stations were located along the coast between the Karatas Cape in the east and Goksu River delta in the west (Fig. 1). Seyhan, Tirtar and Goksu stations were directly influenced by the rivers of (Tarsus and Seyban), Lamas and Goksu respectively.

Materials in present study were sampled with two different trawler boats of similar characteristics. Samples were collected by deep trawl net, with a cod-end mesh size of 28 mm stretched, towed approximately 1.5 to 4.5 km.

Stations (Fig. 1) were visited for the first time on March 19, 1985 and continued at approximately 9 months until December 20, 1985. Collections were done randomly. The trawled fish were sorted by species, transferred

into the plastic containers of 5 litres and preserved in 10% formalin solution (Healey, 1978; Hesthagen, 1977; Mais, 1972) immediately after separation of the catch.

The definition of 11 morphometric measurements of individual fish were shown in Fig. 2. Morphometrics and sex of a total of 1772 fish were determined and 1542 specimens were used for the discrimination. For the elimination of the differences in percentage ratios of males and females on the results of Discriminant Analyses, at different stations, the percentage of males and females was equalized.

Leiognathus klunzingeri reaches the sexual maturity at rather small size (50 mm), since its maximum length is 120 mm (Bingel, 1987). It is difficult to discriminate this species on their sex due to its small size. Hence the incorrect determination of sex of *L. klunzingeri* would possibly influence the results. Because of higher probability of wrong sex discrimination of smaller individuals, it was decided to analyse larger individuals.

In order to select perfectly matured individuals, time series of the length frequency distribution given in Bingel (1987) was examined. Individuals between 50 and 60 mm in length were considered as the first sexually mature group using Petersen Method, Holden & Raitt (1974). To avoid further errors, Discriminant Analyses were made for 60 mm and longer individuals. Total length ranges of the samples and number of sampled individuals collected from four locations are given in Table 1.

Table 1. Minimum, maximum total length ranges and number of specimens at four sampling stations

Whole data	STATIONS							
	TUZLA		SEYHAN		TIRTAR		GOKSU	
	min	max	min	max	min	max	min	max
ROW DATA								
Total length (mm)	28	96	26	115	41	86	32	98
* of specimens	498		498		278		498	
EQUALIZED DATA								
Total length (mm)	60	96	60	91	60	86	60	94
* of specimens	219		194		131		227	

During washing of each fish in the laboratory, under cold running tap water, the body of each specimen was turned in a tail-to-head direction and the washing operation was done with weakly flowing water. Sub-sampling was performed using the method described by Holden & Raitt (1974). During the length measurement, each individual was laid on its right side, the jaws being closed and the length was measured by means of a clipper as a straight line. The distance between the tips of the clipper was read off in millimetric scale. The weighing was done on an electronic balance (accuracy 0.01 gr).

According to Blackith and Reyment (1971), the weight unit can be used for the standardization of the row data. It is well known that the weight is a power of the length (Ricker, 1975). Based on this idea, the weight unit was used in present study, for the standardization of the row




Fig. 1. Sampling locations in the Mersin Bay [black square]: TUZLA, [white square]: SEYHAN, [hatched square]: TIRTAR, [white square]: GOKSU

data. In order to prevent any bias in total weight measurements due to food factor and different maturation stage, the weight after cleaning was used for standardization. The fish was laid straight on a millimetric paper. Much care was paid to fit the axis, which passes through maxillary symphysis (U) to the membraneous edge of caudal fin at fork (F) to the straight line drawn in a horizontal plane on the millimetric paper (Fig. 2). For dissection through gill-cover notch at an angle of 90° between the above described axis and the gill-cover notch was arranged and dissection was made by one stroke.




Fig. 2. Measured morphometric characteristics of *L. klunzingeri*

TL = Total length, SL = Standard length, DBL = Dorsal body length, UHL = Upper head length, PPL = Pre-Pectoral length, POL = Pre-Orbital length, OD = Orbital diameter, POD = Post-Orbital distance, FL = Fork length, PDL = Pre-Anterior dorsal length, PAL = Pre-Anal length

[Figure is redrawn from (Tortonese 1975)]

The Generalized Distance of Mahalanobis (Mahalanobis et al., (1949) cited in Weber, 1972) was used in the present study for the numerical discrimination of the stocks of silverbelly:

$$D^2 = (b_1 d_1 + b_2 d_2 + b_3 d_3 + \dots + b_m d_m)/m$$

Where:

D^2 : Generalized Mahalanobis Distance

b_i : discriminant functions of the (i)'th measurement
 d_i : mean difference of the (i)'th measurement
 m : number of measured characters

Clustering technique of Davis (1973) was combined with Generalized Distance of Mahalanobis to obtain qualitative relationship between stocks. Significant testing of Sneath & Sokal (1973) was used to observe the presence or absence of significance between the groups for the calculated D^2 values.

RESULTS AND DISCUSSION

Mahalanobis Distance values and calculated significant testing results are shown in four different local areas in Table 2.

Table 2. Generalized Mahalanobis Distances (D^2) of *L. klunzingeri* longer than 60 mm, above the diagonal. Results of the significant testing were shown below the diagonal; (*) significant

STATION	TUZLA D^2	SEYHAN D^2	TIRTAR D^2	GOKSU D^2
TUZLA		0.473	0.475	0.608
SEYHAN	*		0.757	0.857
TIRTAR	*	*		0.429
GOKSU	*	*	*	

As can be seen from the Table 2 the two least Generalized Distances were found for Tirtar-Goksu (0.429) and for Tuzla-Seyhan (0.473). The mean of the combinations, formed by the members of this pair of groups, is 1.5 times (0.674) larger than any of two groups. Maximum Generalized Distance value was found to be 0.857 for Seyhan-Goksu. The F values calculated for Silverbellies larger than 60 mm using the methods described by Davis (1973), Overall & Klett (1972) and Sneath & Sokal (1973) at 5% error possibility were 3.22, 2.15, 3.28, 3.66, 7.08, 2.12 for Tuzla-Seyhan, Tuzla-Tirtar, Tuzla-Goksu, Seyhan-Tirtar, Seyhan-Goksu, and Tirtar-Goksu respectively. These calculated F values were much higher than the F values read from the table of 1.83, 1.83, 1.82, 1.83, 1.82 and 1.82 respectively. Therefore the hypothesis, that »there is no difference between the groups« was rejected and concluded that four different stocks of silverbellies (Tuzla, Seyhan, Tirtar and Goksu) inhabit the Bay of Mersin.

L. klunzingeri move towards the estuaries even into the mouth of rivers to lay their eggs (Aksiray, 1954). Possible niches for the newly hatched larvae can be either the shallow estuaries or the mouth of rivers. Tarsus and Seyhan rivers flow into the northeastern part of the Mersin Bay, whereas Lamas and Goksu rivers to southwestern parts (Fig. 1). The mouths of Seyhan and Tarsus rivers may be the habitats for Tuzla and Seyhan stocks and the mouths of Lamas and Goksu rivers for the Tirtar and Goksu stocks, respectively.

Results of the cluster analysis (Fig. 3) support also the existence of four different stocks mentioned above. However, the dissimilarity (D^2 value) bet-

ween the samples of Seyhan and Tuzla was higher (0.47) than the samples of Tirtar and Goksu (0.43). As can be seen from the Fig. 3, Tuzla-Seyhan stocks and Tirtar-Goksu stocks differ from each other with a Generalized Distance value of 0.7.




Fig. 3. Dendrogram of Generalized Distance matrix of *L. klunzingeri*

Individuals of Tirtar and Goksu stocks show much more similarity among themselves than Tuzla and Seyhan stock do. Here, Deli Cape seems to be a barrier in the genetic exchange between the individuals of Tuzla and Seyhan stocks. This situation would result in less similarity between the individuals of Tuzla and Seyhan stocks than Tirtar and Goksu stocks.

CONCLUSION

The stock composition of the *Leiognathus klunzingeri*, collected from the northeastern Mediterranean (Mersin Bay) was studied statistically employing the Generalized Distance of Mahalanobis.

Four different stocks of the *L. klunzingeri* were distinguished, i.e., Tuzla, Seyhan, Tirtar and Goksu stocks. This characteristic has been explained by utilizing the numerical compositions of the generalized distances (Table 2) for related stations.

Results of the Cluster analysis also supported the existence of four different stocks. The individuals of Tirtar and Goksu stocks show more similarity than individuals of Tuzla and Seyhan stocks (Fig. 3).

REFERENCES

- Aksiray, F. 1954. Türkiye deniz balıkları tayin anahtarı, I. U. Fen Fak. Hidrobiyol. Arast. Enst. yay. Sayı, pp 1-277, Pulhan matbaası 1st. 283 pp.
- Ben-Tuvia, A. 1966. Red Sea fishes recently found in the Mediterranean. Copeia, 2: 254-275.
- Bingel, F. 1981. Erdemli-ICEL bölgesi balıkçılığı geliştirme projesi kesin raporu. ODTU DBE Erdemli — ICEL. Proje No: 80.07.00-10, 154 pp.
- Bingel, F. 1987. Dogu Akdeniz'de kiya balıkçılığı av alanlarında sayısal balıkçılık projesi kesin raporu. (Quantitative fishery on the fishing grounds of the coastal fishery in the eastern Mediterranean-Final report). Proje No: 80070011, IMS-METU, 312 pp.
- Blackith, R. E. & R. A. Reyment. 1971. Multivariate morphometric. Academic Press, Inc; New York, 412 pp.
- Davis, J. C. 1973. Statistics and Data Analysis in Geology. John Wiley and Sons, inc. USA, 550 pp.
- Ehrich, S., U. Rempe. 1980. Morphometric discrimination between hake populations (Osteichthyes, Gadiformes, Gn. *Merluccius*) from the Northeast Pacific by use of a size-independent discriminant analysis. Mitteilungen aus dem Inst. fuer Seefisch., der Bundesforschungsanstalt fuer Fischerei, Hamburg, 31: 50 pp.
- Erazi, R. A. 1943. *Leiognathus mediterraneus* nov. sp. Compte Rend. Soc. Turq. Sci. Phys., 10: 49-53.
- Healey, M. C. 1978. Fecundity changes in exploited populations of Lake Whitefish (*Coregonus clupeaformis*) and Lake Trout (*Salvelinus namaycush*). Vol. 35: 945-950.
- Hesthagen, I. H. 1977. Migrations, breeding, and growth in *Pomatoschistus minutus* (PALLAS) (Pisces, Gobiidae) in Oslofjorden. Sarsia, 63: 17-26.
- Holden, M. J., D. F. S. Raitt, (Eds.). 1974. Manual of fisheries science. Part 2 — Methods of resource investigation and their application, FAO Fish. Tech. Rep: (115) Rev. 1: 214 pp.
- Kosswig, C. 1974. Modificability a neglected factor for area expansion in marine fish. Istanbul Univ. Fen Fak. Mecm, Ser. B, 39 (1-2): 1-7.
- Mahalanobis, P. C., D. N. Majumda & R. C. Rao. 1949. Anthropometric survey of the united provinces. A statistical study. Sankhya, Calcutta 9: 89-324.
- Mais, K. F. 1972. A subpopulation study of the Pacific sardine. Calif. Fish Game, 58: 296-314.
- Overall, D., C. J. Klett. 1972. Applied multivariate analysis. Mc Graw-Hill Book Company, New York, 500 pp.
- Ricker, H. E. 1975. Computation and Interpretation of Biological statistics of fish populations. Bull. Fish. Res. Board Can., 191, 382 pp.
- Sneath, P. H. A., R. R. Sokal. 1973. Numerical taxonomy. The principles and practice of numerical classification. Freeman Co. San Francisco, 573 pp.
- Steindachner, F. 1898. Ueber einige neue Fischarten aus dem Roten Meere, gesammelt waehrend der I and II. Oesterreichischen Expedition nach dem Rothen Meere in den Jahren 1895-1896 und 1897-1898. S. B. Akad. Wiss. Wien 107, 1 Abt. 780-788.
- Tortonese, E. 1975. Fauna d' Italia, Vol. XI. Osteichthyes pesci ossei, pp. 496-503. Officine Grafiche Calderini, Bologna, 636 pp.
- Weber, E. 1972. Grundriss der biologischen Statistik. 7. Aufl. VEB Gustav Fisher Verlag Jena, 706 pp.

PRIMJENA MAHALANOBIS-OVE FUNKCIJE DISKRIMINACIJE
NA ODREĐIVANJE MORFOMETRIJSKIH RAZLIKA IZMEĐU STOKOVA
LEIOGNATHUS KLUNZINGERI STEINDACHNER U ZALJEVU MERSIN

Dursun Avsar, Ferit Bingel i Mustafa Ünsal

Institut za istraživanje mora, M. E. T. U., Erdemly, Icel, Turska

KRATKI SADRŽAJ

Diferencijacija stocka *Leiognathus klunzingeri* sa četiri lokaliteta sjeveroistočnog Mediterana izučavana je na osnovu morfometrije. Primijenjena je Mahalanobis-ova funkcija diskriminacije i ustanovljeno da postoje četiri različita stoka *Leiognathus klunzingeri* Steindachner u zaljevu Mersin.